

Study of the finite-temperature physics of a lattice SU(4) gauge BSM model.

Venkitesh Ayyar¹ Thomas Degrand¹ Daniel Hackett¹
William Jay¹ Ethan Neil^{1,3} Benjamin Svetitsky² Yigal
Shamir².

¹University of Colorado, Boulder, ²Tel Aviv University, ³RIKEN-BNL Research Center.



University of Colorado
Boulder



Mon, June 19, Lattice 2017, Granada, Spain

Work supported by grants from the DOE.

Computational work done using resources provided by Fermilab and local Janus cluster.



University of Colorado
Boulder

Motivation


- ▶ Study of $SU(4)$ gauge theory with fermions in multiple representations in $(3+1)D$.
- ▶ Interesting in the context of partial compositeness^{1,2}(W. Jay's talk).
- ▶ Looking for **confinement** and **chiral** transitions.
- ▶ Can have different transitions for each representation.

¹D. B. Kaplan and H. Georgi, Phys. Lett. B 136, 183 (1984).

²G. Ferretti, JHEP 06 (2014), 142.

Model details

- ▶ SU(4) gauge theory.
- ▶ NDS action³(nHYP Dislocation Suppression).
- ▶ Clover improved Wilson fermions.
- ▶ 2 flavors of fermions in the fundamental representation(F).
- ▶ 2 flavors of fermions in two-index anti-symmetric representation (A_2)
- ▶ 3D bare parameter space : $\beta, \kappa_4, \kappa_6$.
- ▶ Finite temperature study.



³DeGrand, Shamir, Svetitsky 2014

Types of transitions (... for this talk)

Confinement transition

- ▶ Low temperatures \rightarrow color singlets. No individual quarks.
- ▶ High temperatures \rightarrow Individual quarks can exist.
- ▶ Transition line moves with N_t .

Bulk Transition

- ▶ Unphysical lattice transition.
- ▶ Transition line at fixed bare parameters is independent of N_t .

Polyakov loop expectation value

In pure gauge theory

- ▶ $\langle P \rangle$ order parameter for SSB of centre symmetry $Z(4)$.
- ▶ Confined $\implies \langle P \rangle = 0$.
- ▶ Deconfined, $\implies \langle P \rangle \neq 0$.

Full multi-rep theory


- ▶ F fermions break $Z(4)$ completely.
- ▶ A_2 fermions break $Z(4)$ to $Z(2)$.
- ▶ Pseudo-order parameter.
- ▶ $\langle P \rangle = e^{-\beta F}$
- ▶ $P^F \equiv e^{-\text{Free energy of F quark}}$.
- ▶ Similarly, can define $P^{A_2} \equiv e^{-\text{Free energy of } A_2 \text{ quark}}$.
- ▶ Can look at confinement-deconfinement for each representation.

Flow anisotropy

- ▶ Wilson flow observable $\langle t^2 E(t) \rangle$ can be used to diagnose the phase⁴.
- ▶ $\langle t^2 E(t) \rangle = \langle t^2 E_{ss}(t) \rangle + \langle t^2 E_{st}(t) \rangle$
- ▶ Spatial and temporal values deviate in deconfined phase.
- ▶ In previous work, $t^2 \langle E_{ss}(t) - E_{st}(t) \rangle$.
- ▶ We use $\langle \frac{E_{ss}(t)}{E_{st}(t)} \rangle - 1$.

⁴ S. Datta, S. Gupta, and A. Lytle, Phys. Rev. D94, 094502 (2016);

M. Wandelt, F. Knechtli, and M. Gunther, JHEP 10, 061 (2016)




Lattice simulation details

- ▶ Finite temperature study.
- ▶ Lattice sizes $12^3 \times 6$ and $16^3 \times 8$.
- ▶ Both diagnostics (PL and flow) are consistent.
- ▶ Empirically determine a cut to distinguish confined and deconfined phases
- ▶ For Polyakov loop, the variance of $\arg(P) < 1.0$.
- ▶ 30-200 independent configs for each ensemble.
- ▶ Around 500 ensembles.

Sextet-only limit

Previous work⁵ with A_2
fermions (different action)




- ▶ Bulk transition followed by
- ▶ Confinement transition
- ▶ $\kappa_{critical}$

⁵T. DeGrand, Y. Liu, E. Neil, Y. Shamir, B. Svetitsky, PRD 91, (2015).


Sextet-only limit

Previous work⁵ with A_2 fermions (different action)





- ▶ Bulk transition followed by
- ▶ Confinement transition
- ▶ $\kappa_{critical}$.

NDS action in the A_2 -only limit. ($12^3 \times 6$ lattice)





- ▶ Confinement transition
- ▶ $\kappa_{critical}$.

⁵T. DeGrand, Y. Liu, E. Neil, Y. Shamir, B. Svetitsky, PRD 91, (2015).




$$\beta = 9.6$$



No Bulk transition in region of interest with NDS action.


Phase diagram

Simulation with both representations:



Variation with temperature

- Curve shifts to the right as temperature $T = \frac{1}{aN_t}$ decreases.



No phase separation for different reps

- ▶ Polyakov loop for F and A_2 at fixed β and κ_4 .
- ▶ All observables respond simultaneously.
- ▶ Same in entire parameter space.
- ▶ **Single transition.**
- ▶ Transition seems first order.





Conclusions

- ▶ Finite temperature study of $SU(4)$ gauge theory with 2 fermion reps.
- ▶ NDS action helps avoid the bulk transition.
- ▶ Confinement transitions for both reps coincide.
- ▶ Chiral transition for both reps? (D. Hackett's talk)

THANK YOU

Order of transition



- Polyakov loop and Plaquette discontinuous \implies 1st order confinement transition.

Higher-rep Polyakov loop



- ▶ $\Omega_x^F = \prod_t U_{x,t}^F$
- ▶ $P^F = \text{Tr} \Omega^F$

- ▶ $\Omega_x^{A_2} = \prod_t U_{x,t}^{A_2}$
- ▶ $P^{A_2} = \text{Tr} \Omega^{A_2}$

- ▶ $U^{A_2} = \frac{1}{2} \left[U_{ij}^F U_{kl}^F - U_{il}^F U_{kj}^F \right]$
- ▶ $P^{A_2} = \frac{1}{2} \left[(\text{Tr} \Omega^F)^2 - \text{Tr}(\Omega^F)^2 \right]$

Comparison for $\beta = 7.75$ and $\beta = 7.4$.

- Curve shifts to the right.



Phase diagram comparison

- ▶ Comparing the confinement transition for F vs A_2 .
- ▶ Both transitions co-incide.

