

Scattering from finite-volume energies including higher partial waves and multiple decay channels

Colin Morningstar

Carnegie Mellon University

35th International Symposium
on Lattice Field Theory

Granada, Spain

June 19, 2017



Introduction

- purpose: provide practitioner's guide plus software for carrying out two-particle scattering studies in lattice QCD
- recast in terms of K -matrix and a Hermitian "box matrix" $B^{(P)}$
- provide explicit box matrix elements in block diagonal basis
 - several total momenta
 - total spins $S \leq 2$
 - orbital angular momenta $L \leq 6$
- software to include higher partial waves, multi-channels
- discuss two fitting strategies

- collaborators: John Bulava, Ben Hörz, Bijit Singha, Jacob Fallica, Drew Hanlon, Ruairi Brett
- related talks:
 - John Bulava: Pion-pion scattering 17:20 today
 - Ben Hörz: Multi-hadron spectroscopy 15:30 Thursday

Scattering phase shifts in lattice QCD timeline

- DeWitt 1956: finite-volume energies related to scattering phase shifts
- Lüscher 1986: fields in a cubic box
- Rummukainen and Gottlieb 1995: nonzero total momenta
- Kim, Sachrajda, and Sharpe 2005: derivation reworked
- explosion of papers since then
- Briceno 2014: generalized to arbitrary spin, multiple channels

Two-particle correlator in finite-volume

- correlator of two-particle operator σ in finite volume

$$C^L(P) = \begin{array}{c} \text{Diagram 1} + \text{Diagram 2} \\ + \text{Diagram 3} + \dots \end{array}$$

The diagrams show a series of terms in the expansion of the correlator $C^L(P)$. Each term consists of a chain of circles connected by blue lines. The first and last circles are labeled σ and σ^\dagger respectively. The intermediate circles are labeled iK . Dashed boxes enclose the first and last circles of each term, along with the adjacent iK circles. The first term has one iK circle. The second term has two iK circles. The third term has three iK circles. Ellipses indicate the series continues.

- Bethe-Salpeter kernel

$$\text{Diagram} = \begin{array}{c} \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} \\ + \text{Diagram 4} + \text{Diagram 5} \end{array}$$

The diagram shows the decomposition of the Bethe-Salpeter kernel iK into five terms. The first term is a simple crossing of two lines. The second and third terms are diagrams with two internal lines and two vertices (black and blue dots). The fourth and fifth terms are diagrams with a single internal line and two vertices (green dots).

- $C^\infty(P)$ has branch cuts where two-particle thresholds begin
- momentum quantization in finite volume: cuts \rightarrow series of poles
- C^L poles: two-particle energy spectrum of finite volume theory

Corrections from finite momentum sums

- finite-volume momentum sum is infinite-volume integral plus correction \mathcal{F}

$$\boxed{\text{Diagram}} = \text{Diagram} + \mathcal{F}$$

- define the following quantities: $A, A', i\mathcal{M}$, invariant scattering amplitude $i\mathcal{M}$

$$\begin{aligned} A &= \sigma + \sigma \text{ (with } iK \text{)} + \dots \\ A' &= \sigma' + iK \text{ (with } \sigma' \text{)} + \dots \\ i\mathcal{M} &= iK + iK \text{ (with } iK \text{)} + \dots \end{aligned}$$

Quantization condition

- subtracted correlator $C_{\text{sub}}(P) = C^L(P) - C^\infty(P)$ given by

$$C_{\text{sub}}(P) = \begin{array}{c} \textcircled{A} \text{---} \mathcal{F} \text{---} \textcircled{A'} + \textcircled{A} \text{---} \mathcal{F} \text{---} \textcircled{iM} \text{---} \mathcal{F} \text{---} \textcircled{A'} \\ + \textcircled{A} \text{---} \mathcal{F} \text{---} \textcircled{iM} \text{---} \mathcal{F} \text{---} \textcircled{iM} \text{---} \mathcal{F} \text{---} \textcircled{A'} + \dots \end{array}$$

- sum geometric series

$$C_{\text{sub}}(P) = A \mathcal{F} (1 - iM\mathcal{F})^{-1} A'$$

- poles of $C_{\text{sub}}(P)$ are poles of $C^L(P)$ from $\det(1 - iM\mathcal{F}) = 0$
- key tool: for $g_c(\mathbf{p})$ spatially contained and regular

$$\frac{1}{L^3} \sum_{\mathbf{p}} g_c(\mathbf{p}) = \int \frac{d^3k}{(2\pi)^3} g_c(\mathbf{k}) + O(e^{-mL})$$

$$\frac{1}{L^3} \sum_{\mathbf{p}} \frac{g_c(\mathbf{p}^2)}{(\mathbf{p}^2 - a^2)} = \frac{1}{L^3} \sum_{\mathbf{p}} \frac{g_c(a^2)}{(\mathbf{p}^2 - a^2)} + \int \frac{d^3k}{(2\pi)^3} \frac{g_c(\mathbf{p}^2) - g_c(a^2)}{(\mathbf{p}^2 - a^2)} + O(e^{-mL})$$

Kinematics

- work in spatial L^3 volume with periodic b.c.
- total momentum $\mathbf{P} = (2\pi/L)\mathbf{d}$, where \mathbf{d} vector of integers
- calculate lab-frame energy E of two-particle interacting state in lattice QCD
- boost to center-of-mass frame by defining:

$$E_{\text{cm}} = \sqrt{E^2 - \mathbf{P}^2}, \quad \gamma = \frac{E}{E_{\text{cm}}},$$

- assume N_d channels
- particle masses m_{1a}, m_{2a} and spins s_{1a}, s_{2a} of particle 1 and 2
- for each channel, can calculate

$$\mathbf{q}_{\text{cm},a}^2 = \frac{1}{4}E_{\text{cm}}^2 - \frac{1}{2}(m_{1a}^2 + m_{2a}^2) + \frac{(m_{1a}^2 - m_{2a}^2)^2}{4E_{\text{cm}}^2},$$
$$u_a^2 = \frac{L^2 \mathbf{q}_{\text{cm},a}^2}{(2\pi)^2}, \quad s_a = \left(1 + \frac{(m_{1a}^2 - m_{2a}^2)}{E_{\text{cm}}^2}\right) \mathbf{d}$$

Quantization condition re-expressed

- E related to S matrix (and phase shifts) by

$$\det[1 + F^{(P)}(S - 1)] = 0$$

- F matrix in $JLSa$ basis states given by

$$\begin{aligned} \langle J' m_{J'} L' S' a' | F^{(P)} | J m_J L S a \rangle = & \delta_{a'a} \delta_{S'S} \frac{1}{2} \left\{ \delta_{J'J} \delta_{m_{J'} m_J} \delta_{L'L} \right. \\ & \left. + \langle J' m_{J'} | L' m_{L'} S m_S \rangle \langle L m_L S m_S | J m_J \rangle W_{L' m_{L'}; L m_L}^{(Pa)} \right\} \end{aligned}$$

- total ang mom J, J' , orbital L, L' , spin S, S' , channels a, a'
- W given by

$$\begin{aligned} -i W_{L' m_{L'}; L m_L}^{(Pa)} = & \sum_{l=|L'-L|}^{L'+L} \sum_{m=-l}^l \frac{\mathcal{Z}_{lm}(s_a, \gamma, u_a^2)}{\pi^{3/2} \gamma u_a^{l+1}} \sqrt{\frac{(2L'+1)(2l+1)}{(2L+1)}} \\ & \times \langle L' 0, l 0 | L 0 \rangle \langle L' m_{L'}, l m | L m_L \rangle. \end{aligned}$$

- above expressions apply for both distinguishable and indistinguishable particles

RGL shifted zeta functions

- compute Rummukainen-Gottlieb-Lüscher (RGL) shifted zeta functions \mathcal{Z}_{lm} using

$$\begin{aligned}\mathcal{Z}_{lm}(s, \gamma, u^2) &= \sum_{\mathbf{n} \in \mathbb{Z}^3} \frac{\mathcal{Y}_{lm}(\mathbf{z})}{(\mathbf{z}^2 - u^2)} e^{-\Lambda(\mathbf{z}^2 - u^2)} + \delta_{l0} \frac{\gamma\pi}{\sqrt{\Lambda}} F_0(\Lambda u^2) \\ &+ \frac{i^l \gamma}{\Lambda^{l+1/2}} \int_0^1 dt \left(\frac{\pi}{t}\right)^{l+3/2} e^{\Lambda t u^2} \sum_{\substack{\mathbf{n} \in \mathbb{Z}^3 \\ \mathbf{n} \neq 0}} e^{\pi i \mathbf{n} \cdot \mathbf{s}} \mathcal{Y}_{lm}(\mathbf{w}) e^{-\pi^2 \mathbf{w}^2 / (t\Lambda)}\end{aligned}$$

- where

$$\mathbf{z} = \mathbf{n} - \gamma^{-1} \left[\frac{1}{2} + (\gamma - 1) s^{-2} \mathbf{n} \cdot \mathbf{s} \right] \mathbf{s},$$

$$\mathbf{w} = \mathbf{n} - (1 - \gamma) s^{-2} \mathbf{s} \cdot \mathbf{n} \mathbf{s}, \quad \mathcal{Y}_{lm}(\mathbf{x}) = |\mathbf{x}|^l Y_{lm}(\hat{\mathbf{x}})$$

$$F_0(x) = -1 + \frac{1}{2} \int_0^1 dt \frac{e^{tx} - 1}{t^{3/2}}$$

- choose $\Lambda \approx 1$ for convergence of the summation
- integral done Gauss-Legendre quadrature
- $F_0(x)$ given in terms of Dawson or erf function

K matrix

- quantization condition relates single energy E to entire S -matrix
- cannot solve for S -matrix (except single channel, single wave)
- approximate S -matrix with functions depending on handful of fit parameters
- obtain estimates of fit parameters using many energies
- easier to parametrize Hermitian matrix than unitary matrix
- introduce K -matrix (Wigner 1946)

$$S = (1 + iK)(1 - iK)^{-1} = (1 - iK)^{-1}(1 + iK)$$

- Hermiticity of K -matrix ensures unitarity of S -matrix
- with time reversal invariance, K -matrix must be real and symmetric

K matrix

- rotational invariance implies

$$\langle J' m_{J'} L' S' a' | K | J m_J L S a \rangle = \delta_{J' J} \delta_{m_{J'} m_J} K_{L' S' a'; L S a}^{(J)}(E)$$

where $K^{(J)}$ is real, symmetric, independent of m_J

- invariance under parity gives

$$K_{L' S' a'; L S a}^{(J)}(E) = 0 \quad \text{when} \quad \eta_{1a'}^{P'} \eta_{1a}^P \eta_{2a'}^{P'} \eta_{2a}^P (-1)^{L'+L} = -1,$$

where η_{ja}^P is intrinsic parity of particle j in channel a

- multichannel effective range expansion (Ross 1961)

$$K_{L' S' a'; L S a}^{-1}(E) = q_{a'}^{-L'-\frac{1}{2}} \widehat{K}_{L' S' a'; L S a}^{-1}(E_{\text{cm}}) q_a^{-L-\frac{1}{2}},$$

where $\widehat{K}_{L' S' a'; L S a}^{-1}(E_{\text{cm}})$ real, symmetric, analytic function of E_{cm}

The “box matrix” B

- effective range expansion suggests writing

$$K_{L'S'a'; L_S a}^{-1}(E) = u_{a'}^{-L' - \frac{1}{2}} \tilde{K}_{L'S'a'; L_S a}^{-1}(E_{\text{cm}}) u_a^{-L - \frac{1}{2}}$$

since $\tilde{K}_{L'S'a'; L_S a}^{-1}(E_{\text{cm}})$ behaves smoothly with E_{cm}

- quantization condition can be written

$$\det(1 - B^{(P)} \tilde{K}) = \det(1 - \tilde{K} B^{(P)}) = 0$$

- we define the **box matrix** by

$$\begin{aligned} \langle J' m_{J'} L' S' a' | B^{(P)} | J m_J L S a \rangle &= -i \delta_{a' a} \delta_{S' S} u_a^{L' + L + 1} W_{L' m_{L'}; L m_L}^{(Pa)} \\ &\times \langle J' m_{J'} | L' m_{L'}, S m_S \rangle \langle L m_L, S m_S | J m_J \rangle \end{aligned}$$

- box matrix is **Hermitian** for u_a^2 real
- quantization condition can also be expressed as

$$\det(\tilde{K}^{-1} - B^{(P)}) = 0$$

- these determinants are **real**

Block diagonalization

- quantization condition involves determinant of infinite matrix
- make practical by (a) transforming to a block-diagonal basis and (b) truncating in orbital angular momentum
- for symmetry operation G , define unitary matrix

$$\langle J' m_{J'} L' S' a' | Q^{(G)} | J m_J L S a \rangle = \begin{cases} \delta_{J' J} \delta_{L' L} \delta_{S' S} \delta_{a' a} D_{m_{J'} m_J}^{(J)}(R), & (G = R), \\ \delta_{J' J} \delta_{m_{J'} m_J} \delta_{L' L} \delta_{S' S} \delta_{a' a} (-1)^L, & (G = I_s), \end{cases}$$

where $D_{m' m}^{(J)}(R)$ Wigner rotation matrices, R ordinary rotation, I_s spatial inversion

- can show that box matrix satisfies

$$B^{(GP)} = Q^{(G)} B^{(P)} Q^{(G)\dagger}.$$

- if G in little group of P , then $GP = P$, $Gs_a = s_a$ and

$$[B^{(P)}, Q^{(G)}] = 0, \quad (G \text{ in little group of } P).$$

- can use eigenvectors of $Q^{(G)}$ to block diagonalize $B^{(P)}$

Block diagonalization (con't)

- block-diagonal basis

$$|\Lambda\lambda nJLSa\rangle = \sum_{m_J} c_{m_J}^{J(-1)^L; \Lambda\lambda n} |Jm_JLSa\rangle$$

- little group irrep Λ , irrep row λ , occurrence index n
- transformation coefficients depend on J and $(-1)^L$, not on S, a
- replaces m_J by (Λ, λ, n)
- group theoretical projections with Gram-Schmidt used to obtain coefficients
- use notation and irrep matrices from PRD 88, 014511 (2013)

Block diagonal basis

- $|m_J\rangle$ abbreviates $|Jm_JLSa\rangle$ with parity $\eta = (-1)^L$ for $P = 0$

Λ	λ	J^η	n	Basis vectors
$A_{1\eta}$	1	0^η	1	$ 0\rangle$
$G_{1\eta}$	1	$\frac{1}{2}^\eta$	1	$ \frac{1}{2}\rangle$
$G_{1\eta}$	2	$\frac{1}{2}^\eta$	1	$ \frac{1}{2}\rangle$
$T_{1\eta}$	1	1^η	1	$\frac{1}{\sqrt{2}}(1\rangle - -1\rangle)$
$T_{1\eta}$	2	1^η	1	$\frac{-i}{\sqrt{2}}(1\rangle + -1\rangle)$
$T_{1\eta}$	3	1^η	1	$ 0\rangle$
H_η	1	$\frac{3}{2}^\eta$	1	$ \frac{3}{2}\rangle$
H_η	2	$\frac{3}{2}^\eta$	1	$ \frac{3}{2}\rangle$
H_η	3	$\frac{3}{2}^\eta$	1	$ \frac{3}{2}\rangle$
H_η	4	$\frac{3}{2}^\eta$	1	$ \frac{3}{2}\rangle$
E_η	1	2^η	1	$\frac{1}{\sqrt{2}}(2\rangle + -2\rangle)$
E_η	2	2^η	1	$ 0\rangle$
$T_{2\eta}$	1	2^η	1	$\frac{1}{\sqrt{2}}(1\rangle + -1\rangle)$
$T_{2\eta}$	2	2^η	1	$\frac{i}{\sqrt{2}}(1\rangle - -1\rangle)$
$T_{2\eta}$	3	2^η	1	$\frac{1}{\sqrt{2}}(- 2\rangle + -2\rangle)$
$G_{2\eta}$	1	$\frac{5}{2}^\eta$	1	$\frac{1}{\sqrt{6}}(\frac{5}{2}\rangle - \sqrt{5} \frac{3}{2}\rangle)$
$G_{2\eta}$	2	$\frac{5}{2}^\eta$	1	$\frac{1}{\sqrt{6}}(-\sqrt{5} \frac{3}{2}\rangle + \frac{5}{2}\rangle)$
H_η	1	$\frac{5}{2}^\eta$	1	$\frac{1}{\sqrt{6}}(\frac{3}{2}\rangle + \sqrt{5} \frac{5}{2}\rangle)$
H_η	2	$\frac{5}{2}^\eta$	1	$ \frac{1}{2}\rangle$
H_η	3	$\frac{5}{2}^\eta$	1	$ \frac{1}{2}\rangle$
H_η	4	$\frac{5}{2}^\eta$	1	$\frac{-i}{\sqrt{6}}(\sqrt{5} \frac{5}{2}\rangle + \frac{3}{2}\rangle)$

Block diagonal basis

Λ	λ	J^η	n	Basis vectors $\mathbf{P} = 0$
$A_{2\eta}$	1	3^η	1	$\frac{1}{\sqrt{2}} (2\rangle - -2\rangle)$
$T_{1\eta}$	1	3^η	1	$\frac{1}{4} (\sqrt{5} 3\rangle - \sqrt{3} 1\rangle + \sqrt{3} -1\rangle - \sqrt{5} -3\rangle)$
$T_{1\eta}$	2	3^η	1	$\frac{1}{4} (\sqrt{5} 3\rangle + \sqrt{3} 1\rangle + \sqrt{3} -1\rangle + \sqrt{5} -3\rangle)$
$T_{1\eta}$	3	3^η	1	$ 0\rangle$
$T_{2\eta}$	1	3^η	1	$\frac{1}{4} (\sqrt{3} 3\rangle + \sqrt{5} 1\rangle - \sqrt{5} -1\rangle - \sqrt{3} -3\rangle)$
$T_{2\eta}$	2	3^η	1	$\frac{1}{4} (-\sqrt{3} 3\rangle + \sqrt{5} 1\rangle + \sqrt{5} -1\rangle - \sqrt{3} -3\rangle)$
$T_{2\eta}$	3	3^η	1	$\frac{1}{\sqrt{2}} (2\rangle + -2\rangle)$
$G_{1\eta}$	1	$\frac{7}{2}\eta$	1	$\frac{1}{2\sqrt{3}} (\sqrt{7} \frac{1}{2}\rangle + \sqrt{5} -\frac{7}{2}\rangle)$
$G_{1\eta}$	2	$\frac{7}{2}\eta$	1	$\frac{-1}{2\sqrt{3}} (\sqrt{5} \frac{7}{2}\rangle + \sqrt{7} -\frac{1}{2}\rangle)$
$G_{2\eta}$	1	$\frac{7}{2}\eta$	1	$\frac{1}{2} (\sqrt{3} \frac{5}{2}\rangle - -\frac{3}{2}\rangle)$
$G_{2\eta}$	2	$\frac{7}{2}\eta$	1	$\frac{1}{2} (\frac{3}{2}\rangle - \sqrt{3} -\frac{5}{2}\rangle)$
H_η	1	$\frac{7}{2}\eta$	1	$\frac{1}{2} (\sqrt{3} \frac{3}{2}\rangle + -\frac{7}{2}\rangle)$
H_η	2	$\frac{7}{2}\eta$	1	$\frac{1}{2\sqrt{3}} (-\sqrt{5} \frac{1}{2}\rangle + \sqrt{7} -\frac{7}{2}\rangle)$
H_η	3	$\frac{7}{2}\eta$	1	$\frac{1}{2\sqrt{3}} (\sqrt{7} \frac{7}{2}\rangle - \sqrt{5} -\frac{1}{2}\rangle)$
H_η	4	$\frac{7}{2}\eta$	1	$\frac{1}{2} (\frac{5}{2}\rangle + \sqrt{3} -\frac{3}{2}\rangle)$

Block diagonal basis

Λ	λ	J^η	n	Basis vectors $P = 0$
$A_{1\eta}$	1	4^η	1	$\frac{1}{2\sqrt{6}}(\sqrt{5} 4\rangle + \sqrt{14} 0\rangle + \sqrt{5} -4\rangle)$
E_η	1	4^η	1	$\frac{1}{\sqrt{2}}(2\rangle + -2\rangle)$
E_η	2	4^η	1	$\frac{1}{2\sqrt{6}}(\sqrt{7} 4\rangle - \sqrt{10} 0\rangle + \sqrt{7} -4\rangle)$
$T_{1\eta}$	1	4^η	1	$\frac{1}{4}(3\rangle + \sqrt{7} 1\rangle + \sqrt{7} -1\rangle + -3\rangle)$
$T_{1\eta}$	2	4^η	1	$\frac{1}{4}(3\rangle - \sqrt{7} 1\rangle + \sqrt{7} -1\rangle - -3\rangle)$
$T_{1\eta}$	3	4^η	1	$\frac{1}{\sqrt{2}}(4\rangle - -4\rangle)$
$T_{2\eta}$	1	4^η	1	$\frac{1}{4}(\sqrt{7} 3\rangle - 1\rangle - -1\rangle + \sqrt{7} -3\rangle)$
$T_{2\eta}$	2	4^η	1	$\frac{1}{4}(-\sqrt{7} 3\rangle - 1\rangle + -1\rangle + \sqrt{7} -3\rangle)$
$T_{2\eta}$	3	4^η	1	$\frac{1}{\sqrt{2}}(- 2\rangle + -2\rangle)$
$G_{1\eta}$	1	$\frac{9}{2}\eta$	1	$\frac{1}{2\sqrt{6}}(3 \frac{9}{2}\rangle + \sqrt{14} \frac{1}{2}\rangle + -\frac{7}{2}\rangle)$
$G_{1\eta}$	2	$\frac{9}{2}\eta$	1	$\frac{1}{2\sqrt{6}}(\frac{7}{2}\rangle + \sqrt{14} -\frac{1}{2}\rangle + 3 -\frac{9}{2}\rangle)$
H_η	1	$\frac{9}{2}\eta$	1	$ \frac{3}{2}\rangle$
H_η	1	$\frac{9}{2}\eta$	2	$ -\frac{5}{2}\rangle$
H_η	2	$\frac{9}{2}\eta$	1	$\frac{1}{4}(-\sqrt{7} \frac{9}{2}\rangle + \sqrt{2} \frac{1}{2}\rangle + \sqrt{7} - \frac{7}{2}\rangle)$
H_η	2	$\frac{9}{2}\eta$	2	$\frac{-1}{4\sqrt{3}}(3 \frac{9}{2}\rangle - \sqrt{14} \frac{1}{2}\rangle + 5 -\frac{7}{2}\rangle)$
H_η	3	$\frac{9}{2}\eta$	1	$\frac{-1}{4}(\sqrt{7} \frac{7}{2}\rangle + \sqrt{2} -\frac{1}{2}\rangle - \sqrt{7} -\frac{9}{2}\rangle)$
H_η	3	$\frac{9}{2}\eta$	2	$\frac{-1}{4\sqrt{3}}(5 \frac{7}{2}\rangle - \sqrt{14} -\frac{1}{2}\rangle + 3 -\frac{9}{2}\rangle)$
H_η	4	$\frac{9}{2}\eta$	1	$ -\frac{3}{2}\rangle$
H_η	4	$\frac{9}{2}\eta$	2	$ \frac{5}{2}\rangle$

Block diagonal basis

Λ	λ	J^n	n	Basis vectors $P = (0, 0, 1)$
A_1	1	0^+	1	$ 0\rangle$
A_2	1	0^-	1	$ 0\rangle$
G_1	1	$\frac{1}{2}^+$	1	$ \frac{1}{2}\rangle$
G_1	2	$\frac{1}{2}^+$	1	$ \frac{1}{2}\rangle$
G_1	1	$\frac{1}{2}^-$	1	$ \frac{1}{2}\rangle$
G_1	2	$\frac{1}{2}^-$	1	$ \frac{1}{2}\rangle$
A_1	1	1^-	1	$ 0\rangle$
A_2	1	1^+	1	$ 0\rangle$
E	1	1^+	1	$\frac{1}{\sqrt{2}}(1\rangle + -1\rangle)$
E	2	1^+	1	$\frac{i}{\sqrt{2}}(- 1\rangle + -1\rangle)$
E	1	1^-	1	$\frac{1}{\sqrt{2}}(1\rangle - -1\rangle)$
E	2	1^-	1	$\frac{-i}{\sqrt{2}}(1\rangle + -1\rangle)$
G_1	1	$\frac{3}{2}^+$	1	$ \frac{3}{2}\rangle$
G_1	2	$\frac{3}{2}^+$	1	$ \frac{3}{2}\rangle$
G_1	1	$\frac{3}{2}^-$	1	$ \frac{3}{2}\rangle$
G_1	2	$\frac{3}{2}^-$	1	$ \frac{3}{2}\rangle$
G_2	1	$\frac{5}{2}^+$	1	$ \frac{5}{2}\rangle$
G_2	2	$\frac{5}{2}^+$	1	$ \frac{5}{2}\rangle$
G_2	1	$\frac{5}{2}^-$	1	$ \frac{5}{2}\rangle$
G_2	2	$\frac{5}{2}^-$	1	$ \frac{5}{2}\rangle$

Block diagonal basis

- $\nu_1 = \frac{1}{\sqrt{2}}(1 + i)$, $\nu_2 = \frac{1}{2\sqrt{3}}(2 - \sqrt{2} + i(2 + \sqrt{2}))$, $\nu_3 = \frac{1}{\sqrt{3}}(\sqrt{2} + i)$

Λ	λ	J^n	n	Basis vectors $\mathbf{P} = (1, 1, 1)$
A_1	1	3^+	1	$\frac{1}{2\sqrt{6}}(\sqrt{3} 3\rangle + 2\nu_1 2\rangle + i\sqrt{5} 1\rangle - \sqrt{5} -1\rangle + 2\nu_1 -2\rangle - i\sqrt{3} -3\rangle)$
A_1	1	3^-	1	$\frac{1}{2\sqrt{6}}(\sqrt{5} 3\rangle + i\sqrt{3} 1\rangle - 2\sqrt{2}\nu_1^* 0\rangle + \sqrt{3} -1\rangle + i\sqrt{5} -3\rangle)$
A_1	1	3^-	2	$\frac{1}{\sqrt{2}}(- 2\rangle + -2\rangle)$
A_2	1	3^+	1	$\frac{1}{2\sqrt{6}}(\sqrt{5} 3\rangle + i\sqrt{3} 1\rangle - 2\sqrt{2}\nu_1^* 0\rangle + \sqrt{3} -1\rangle + i\sqrt{5} -3\rangle)$
A_2	1	3^+	2	$\frac{1}{\sqrt{2}}(- 2\rangle + -2\rangle)$
A_2	1	3^-	1	$\frac{1}{2\sqrt{6}}(\sqrt{3} 3\rangle + 2\nu_1 2\rangle + i\sqrt{5} 1\rangle - \sqrt{5} -1\rangle + 2\nu_1 -2\rangle - i\sqrt{3} -3\rangle)$
E	1	3^+	1	$\frac{1}{2\sqrt{42}}(7 3\rangle - i\sqrt{15} 1\rangle + 2\sqrt{10}\nu_1^* 0\rangle - \sqrt{15} -1\rangle + 7i -3\rangle)$
E	1	3^+	2	$\frac{-1}{\sqrt{14}}(-2 1\rangle + \sqrt{6}\nu_1 0\rangle + 2i -1\rangle)$
E	2	3^+	1	$\frac{-1}{2\sqrt{14}}(i 3\rangle - 2\sqrt{3}\nu_1^* 2\rangle + \sqrt{15} 1\rangle + i\sqrt{15} -1\rangle - 2\sqrt{3}\nu_1^* -2\rangle + -3\rangle)$
E	2	3^+	2	$\frac{1}{2\sqrt{21}}(-\sqrt{30} 3\rangle + \sqrt{10}\nu_1 2\rangle + i\sqrt{2} 1\rangle - \sqrt{2} -1\rangle + \sqrt{10}\nu_1 -2\rangle + i\sqrt{30} -3\rangle)$
E	1	3^-	1	$\frac{-1}{6\sqrt{2}}(-3\sqrt{3} 3\rangle + 2\nu_1 2\rangle + i\sqrt{5} 1\rangle - \sqrt{5} -1\rangle + 2\nu_1 -2\rangle + 3i\sqrt{3} -3\rangle)$
E	1	3^-	2	$\frac{1}{3\sqrt{2}}(\sqrt{5} 2\rangle - 2\nu_1 1\rangle + 2\nu_1^* -1\rangle + \sqrt{5} -2\rangle)$
E	2	3^-	1	$\frac{-1}{6\sqrt{2}}(i 3\rangle - \sqrt{15} 1\rangle + 2\sqrt{10}\nu_1 0\rangle + i\sqrt{15} -1\rangle - -3\rangle)$
E	2	3^-	2	$\frac{-1}{6}(\sqrt{10}\nu_1 3\rangle + \sqrt{6}\nu_1^* 1\rangle + 2 0\rangle - \sqrt{6}\nu_1 -1\rangle - \sqrt{10}\nu_1^* -3\rangle)$

Box and \tilde{K} matrices in block diagonal basis

- in block-diagonal basis, box matrix has form

$$\langle \Lambda' \lambda' n' J' L' S' a' | B^{(P)} | \Lambda \lambda n J L S a \rangle = \delta_{\Lambda' \Lambda} \delta_{\lambda' \lambda} \delta_{S' S} \delta_{a' a} B_{J' L' n'; J L n}^{(P \Lambda_B S a)}(E)$$

- \tilde{K} -matrix for $(-1)^{L+L'} = 1$ has form

$$\langle \Lambda' \lambda' n' J' L' S' a' | \tilde{K} | \Lambda \lambda n J L S a \rangle = \delta_{\Lambda' \Lambda} \delta_{\lambda' \lambda} \delta_{n' n} \delta_{J' J} \mathcal{K}_{L' S' a'; L S a}^{(J)}(E_{\text{cm}})$$

- $(-1)^{L+L'} = 1 \Rightarrow \eta_{1a'}^{P'} \eta_{2a'}^{P'} = \eta_{1a}^P \eta_{2a}^P$, always applies in QCD
- Λ is irrep for K -matrix, need Λ_B for box matrix
- when $\eta_{1a}^P \eta_{2a}^P = 1$, then $\Lambda_B = \Lambda$

d	LG	Λ_B relationship to Λ when $\eta_{1a}^P \eta_{2a}^P = -1$
$(0, 0, 0)$	O_h	Subscript $g \leftrightarrow u$
$(0, 0, n)$	C_{4v}	$A_1 \leftrightarrow A_2$; $B_1 \leftrightarrow B_2$; E, G_1, G_2 stay same
$(0, n, n)$	C_{2v}	$A_1 \leftrightarrow A_2$; $B_1 \leftrightarrow B_2$; G stays same
(n, n, n)	C_{3v}	$A_1 \leftrightarrow A_2$; $F_1 \leftrightarrow F_2$; E, G stay same

- see PRD 88, 014511 (2013) for notation

K matrix parametrizations

- \tilde{K} matrix in block diagonal basis

$$\langle \Lambda' \lambda' n' J' L' S' a' | \tilde{K} | \Lambda \lambda n J L S a \rangle = \delta_{\Lambda' \Lambda} \delta_{\lambda' \lambda} \delta_{n' n} \delta_{J' J} \mathcal{K}_{L' S' a'; L S a}^{(J)}(E_{\text{cm}})$$

$$\langle \Lambda' \lambda' n' J' L' S' a' | \tilde{K}^{-1} | \Lambda \lambda n J L S a \rangle = \delta_{\Lambda' \Lambda} \delta_{\lambda' \lambda} \delta_{n' n} \delta_{J' J} \mathcal{K}_{L' S' a'; L S a}^{(J)-1}(E_{\text{cm}})$$

- common parametrization

$$\mathcal{K}_{\alpha\beta}^{(J)-1}(E_{\text{cm}}) = \sum_{k=0}^{N_{\alpha\beta}} c_{\alpha\beta}^{(Jk)} E_{\text{cm}}^k$$

- α, β compound indices for (L, S, a)
- another common parametrization

$$\mathcal{K}_{\alpha\beta}^{(J)}(E_{\text{cm}}) = \sum_p \frac{g_{\alpha}^{(Jp)} g_{\beta}^{(Jp)}}{E_{\text{cm}}^2 - m_{j_p}^2} + \sum_k d_{\alpha\beta}^{(Jk)} E_{\text{cm}}^k,$$

- Lorentz invariant form using $E_{\text{cm}} = \sqrt{s}$
- Mandelstam variable $s = (p_1 + p_2)^2$, with p_j four-momentum of particle j

Box matrix elements

- have obtained expressions for $B_{J'L'n'; JLn}^{(\mathbf{P}\Lambda_B S_a)}(E)$ for
- $L \leq 6, S \leq 2$ with $\mathbf{P} = (0, 0, 0), (0, 0, p), p > 0$
- $L \leq 6, S \leq \frac{3}{2}$ with $\mathbf{P} = (0, p, p), (p, p, p), p > 0$
- in tables that follow, we define

R_{lm} is short hand for $(\gamma\pi^{3/2}u_a^{l+1})^{-1}\text{Re } \mathcal{Z}_{lm}(s_a, \gamma, u_a^2)$

I_{lm} is short hand for $(\gamma\pi^{3/2}u_a^{l+1})^{-1}\text{Im } \mathcal{Z}_{lm}(s_a, \gamma, u_a^2)$

Box matrix elements $P = 0, S = 0$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
$\Lambda_B = A_{1g}$						
0	0	1	0	0	1	R_{00}
0	0	1	4	4	1	$\frac{2\sqrt{21}}{7} R_{40}$
0	0	1	6	6	1	$-2\sqrt{2} R_{60}$
4	4	1	4	4	1	$R_{00} + \frac{108}{143} R_{40} + \frac{80\sqrt{13}}{143} R_{60} + \frac{560\sqrt{17}}{2431} R_{80}$
4	4	1	6	6	1	$-\frac{40\sqrt{546}}{1001} R_{40} + \frac{42\sqrt{42}}{187} R_{60} - \frac{224\sqrt{9282}}{46189} R_{80} - \frac{1008\sqrt{26}}{4199} R_{10,0}$
6	6	1	6	6	1	$R_{00} - \frac{126}{187} R_{40} - \frac{160\sqrt{13}}{3553} R_{60} + \frac{840\sqrt{17}}{3553} R_{80} - \frac{2016\sqrt{21}}{7429} R_{10,0}$ $+ \frac{30492}{37145} R_{12,0} - \frac{1848\sqrt{1001}}{37145} R_{12,4}$
$\Lambda_B = A_{2g}$						
6	6	1	6	6	1	$R_{00} + \frac{6}{17} R_{40} - \frac{160\sqrt{13}}{323} R_{60} - \frac{40\sqrt{17}}{323} R_{80} - \frac{2592\sqrt{21}}{7429} R_{10,0}$ $+ \frac{1980}{7429} R_{12,0} + \frac{264\sqrt{1001}}{7429} R_{12,4}$
$\Lambda_B = A_{2u}$						
3	3	1	3	3	1	$R_{00} - \frac{12}{11} R_{40} + \frac{80\sqrt{13}}{143} R_{60}$

Box matrix elements $P = 0, S = 0$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
------	------	------	-----	-----	-----	---------------------

$$\Lambda_B = E_g$$

2	2	1	2	2	1	$R_{00} + \frac{6}{7}R_{40}$
2	2	1	4	4	1	$-\frac{40\sqrt{3}}{77}R_{40} - \frac{30\sqrt{39}}{143}R_{60}$
2	2	1	6	6	1	$\frac{30\sqrt{910}}{1001}R_{40} + \frac{4\sqrt{70}}{55}R_{60} + \frac{8\sqrt{15470}}{1105}R_{80}$
4	4	1	4	4	1	$R_{00} + \frac{108}{1001}R_{40} - \frac{64\sqrt{13}}{143}R_{60} + \frac{392\sqrt{17}}{2431}R_{80}$
4	4	1	6	6	1	$-\frac{8\sqrt{2730}}{1001}R_{40} - \frac{18\sqrt{210}}{187}R_{60} - \frac{128\sqrt{46410}}{46189}R_{80}$ $-\frac{1512\sqrt{130}}{20995}R_{10,0}$
6	6	1	6	6	1	$R_{00} + \frac{114}{187}R_{40} + \frac{480\sqrt{13}}{3553}R_{60} + \frac{280\sqrt{17}}{3553}R_{80} + \frac{1152\sqrt{21}}{7429}R_{10,0}$ $+\frac{30492}{37145}R_{12,0} + \frac{264\sqrt{1001}}{37145}R_{12,4}$

$$\Lambda_B = E_u$$

5	5	1	5	5	1	$R_{00} - \frac{6}{13}R_{40} + \frac{32\sqrt{13}}{221}R_{60} - \frac{672\sqrt{17}}{4199}R_{80} + \frac{1152\sqrt{21}}{4199}R_{10,0}$
---	---	---	---	---	---	--

$$\Lambda_B = T_{1g}$$

4	4	1	4	4	1	$R_{00} + \frac{54}{143}R_{40} - \frac{4\sqrt{13}}{143}R_{60} - \frac{448\sqrt{17}}{2431}R_{80}$
4	4	1	6	6	1	$-\frac{12\sqrt{65}}{143}R_{40} + \frac{42\sqrt{5}}{187}R_{60} + \frac{112\sqrt{1105}}{46189}R_{80} + \frac{576\sqrt{1365}}{20995}R_{10,0}$
6	6	1	6	6	1	$R_{00} - \frac{96}{187}R_{40} - \frac{80\sqrt{13}}{3553}R_{60} + \frac{120\sqrt{17}}{3553}R_{80} + \frac{624\sqrt{21}}{7429}R_{10,0}$ $-\frac{26136}{37145}R_{12,0} + \frac{1584\sqrt{1001}}{37145}R_{12,4}$

Box matrix elements $P = 0, S = 0$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
$\Lambda_B = T_{1u}$						
1	1	1	1	1	1	R_{00}
1	1	1	3	3	1	$\frac{4\sqrt{21}}{21} R_{40}$
1	1	1	5	5	1	$\frac{20\sqrt{3927}}{1309} R_{40} + \frac{4\sqrt{51051}}{2431} R_{60}$
1	1	1	5	5	2	$-\frac{2\sqrt{2805}}{561} R_{40} + \frac{24\sqrt{36465}}{2431} R_{60}$
3	3	1	3	3	1	$R_{00} + \frac{6}{11} R_{40} + \frac{100\sqrt{13}}{429} R_{60}$
3	3	1	5	5	1	$\frac{60\sqrt{187}}{2431} R_{40} + \frac{42\sqrt{2431}}{2431} R_{60} + \frac{112\sqrt{11}}{429} R_{80}$
3	3	1	5	5	2	$\frac{12\sqrt{6545}}{1309} R_{40} - \frac{28\sqrt{85085}}{7293} R_{60}$
5	5	1	5	5	1	$R_{00} + \frac{132}{221} R_{40} + \frac{880\sqrt{13}}{3757} R_{60} + \frac{280\sqrt{17}}{3757} R_{80} + \frac{336\sqrt{21}}{3757} R_{10,0}$
5	5	1	5	5	2	$-\frac{24\sqrt{35}}{1547} R_{40} - \frac{120\sqrt{455}}{3757} R_{60} + \frac{2800\sqrt{595}}{214149} R_{80}$ $+ \frac{88704\sqrt{15}}{356915} R_{10,0}$
5	5	2	5	5	2	$R_{00} - \frac{132}{221} R_{40} + \frac{352\sqrt{13}}{11271} R_{60} + \frac{7056\sqrt{17}}{71383} R_{80}$ $- \frac{12096\sqrt{21}}{71383} R_{10,0}$

Box matrix elements $P = 0, S = 0$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
------	------	------	-----	-----	-----	---------------------

$$\Lambda_B = T_{2g}$$

2	2	1	2	2	1	$R_{00} - \frac{1}{7} R_{40}$
2	2	1	4	4	1	$-\frac{20\sqrt{3}}{77} R_{40} + \frac{40\sqrt{39}}{143} R_{60}$
2	2	1	6	6	1	$\frac{20\sqrt{715}}{1001} R_{40} - \frac{12\sqrt{55}}{55} R_{60} - \frac{32\sqrt{12155}}{36465} R_{80}$
2	2	1	6	6	2	$\frac{190\sqrt{13}}{1001} R_{40} + \frac{8}{11} R_{60} - \frac{32\sqrt{221}}{663} R_{80}$
4	4	1	4	4	1	$R_{00} - \frac{54}{77} R_{40} + \frac{20\sqrt{13}}{143} R_{60}$
4	4	1	6	6	1	$\frac{4\sqrt{2145}}{1001} R_{40} - \frac{2\sqrt{165}}{187} R_{60} - \frac{144\sqrt{36465}}{46189} R_{80} + \frac{384\sqrt{5005}}{20995} R_{10,0}$
4	4	1	6	6	2	$-\frac{60\sqrt{39}}{1001} R_{40} - \frac{124\sqrt{3}}{187} R_{60} + \frac{64\sqrt{663}}{4199} R_{80} + \frac{192\sqrt{91}}{4199} R_{10,0}$
6	6	1	6	6	1	$R_{00} - \frac{32}{119} R_{40} + \frac{80\sqrt{13}}{323} R_{60} - \frac{920\sqrt{17}}{6783} R_{80} - \frac{720\sqrt{21}}{52003} R_{10,0}$ $+ \frac{91608}{260015} R_{12,0} - \frac{5808\sqrt{1001}}{260015} R_{12,4}$
6	6	1	6	6	2	$\frac{40\sqrt{55}}{1309} R_{40} + \frac{120\sqrt{715}}{3553} R_{60} + \frac{80\sqrt{935}}{24871} R_{80} - \frac{4608\sqrt{1155}}{260015} R_{10,0}$ $- \frac{13728\sqrt{55}}{260015} R_{12,0} + \frac{6336\sqrt{455}}{260015} R_{12,4}$
6	6	2	6	6	2	$R_{00} + \frac{632}{1309} R_{40} - \frac{480\sqrt{13}}{3553} R_{60} + \frac{80\sqrt{17}}{6783} R_{80} + \frac{1728\sqrt{21}}{52003} R_{10,0}$ $- \frac{29040}{52003} R_{12,0} - \frac{1056\sqrt{1001}}{52003} R_{12,4}$

$$\Lambda_B = T_{2u}$$

3	3	1	3	3	1	$R_{00} - \frac{2}{11} R_{40} - \frac{60\sqrt{13}}{143} R_{60}$
3	3	1	5	5	1	$-\frac{20\sqrt{11}}{143} R_{40} - \frac{14\sqrt{143}}{143} R_{60} + \frac{112\sqrt{187}}{2431} R_{80}$
5	5	1	5	5	1	$R_{00} + \frac{4}{13} R_{40} - \frac{80\sqrt{13}}{221} R_{60} - \frac{280\sqrt{17}}{4199} R_{80} - \frac{432\sqrt{21}}{4199} R_{10,0}$

Box matrix elements $P = 0$, $S = \frac{1}{2}$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
$\Lambda_B = G_{1g}$						
$\frac{1}{2}$	0	1	$\frac{1}{2}$	0	1	R_{00}
$\frac{1}{2}$	0	1	$\frac{7}{2}$	4	1	$-\frac{4\sqrt{21}}{21} R_{40}$
$\frac{1}{2}$	0	1	$\frac{9}{2}$	4	1	$\frac{2\sqrt{105}}{21} R_{40}$
$\frac{1}{2}$	0	1	$\frac{11}{2}$	6	1	$\frac{4\sqrt{39}}{13} R_{60}$
$\frac{1}{2}$	0	1	$\frac{13}{2}$	6	1	$-\frac{2\sqrt{182}}{13} R_{60}$
$\frac{7}{2}$	4	1	$\frac{7}{2}$	4	1	$R_{00} + \frac{6}{11} R_{40} + \frac{100\sqrt{13}}{429} R_{60}$
$\frac{7}{2}$	4	1	$\frac{9}{2}$	4	1	$-\frac{12\sqrt{5}}{143} R_{40} - \frac{56\sqrt{65}}{429} R_{60} - \frac{224\sqrt{85}}{2431} R_{80}$
$\frac{7}{2}$	4	1	$\frac{11}{2}$	6	1	$-\frac{300\sqrt{7}}{1001} R_{40} + \frac{14\sqrt{91}}{143} R_{60} - \frac{112\sqrt{119}}{7293} R_{80}$
$\frac{7}{2}$	4	1	$\frac{13}{2}$	6	1	$\frac{20\sqrt{6}}{429} R_{40} - \frac{126\sqrt{78}}{2431} R_{60} + \frac{112\sqrt{102}}{4199} R_{80} + \frac{96\sqrt{14}}{323} R_{10,0}$
$\frac{9}{2}$	4	1	$\frac{9}{2}$	4	1	$R_{00} + \frac{84}{143} R_{40} + \frac{128\sqrt{13}}{429} R_{60} + \frac{112\sqrt{17}}{2431} R_{80}$
$\frac{9}{2}$	4	1	$\frac{11}{2}$	6	1	$\frac{24\sqrt{35}}{1001} R_{40} - \frac{56\sqrt{455}}{2431} R_{60} + \frac{1568\sqrt{595}}{138567} R_{80} + \frac{6048\sqrt{15}}{20995} R_{10,0}$
$\frac{9}{2}$	4	1	$\frac{13}{2}$	6	1	$-\frac{64\sqrt{30}}{429} R_{40} + \frac{126\sqrt{390}}{2431} R_{60} - \frac{448\sqrt{510}}{46189} R_{80} - \frac{528\sqrt{70}}{20995} R_{10,0}$
$\frac{11}{2}$	6	1	$\frac{11}{2}$	6	1	$R_{00} - \frac{84}{143} R_{40} - \frac{80\sqrt{13}}{2431} R_{60} + \frac{5880\sqrt{17}}{46189} R_{80}$ $-\frac{336\sqrt{21}}{4199} R_{10,0}$
$\frac{11}{2}$	6	1	$\frac{13}{2}$	6	1	$\frac{30\sqrt{42}}{2431} R_{40} + \frac{80\sqrt{546}}{46189} R_{60} - \frac{720\sqrt{714}}{46189} R_{80} + \frac{55440\sqrt{2}}{96577} R_{10,0}$ $-\frac{4356\sqrt{42}}{37145} R_{12,0} + \frac{1848\sqrt{858}}{37145} R_{12,4}$
$\frac{13}{2}$	6	1	$\frac{13}{2}$	6	1	$R_{00} - \frac{1458}{2431} R_{40} - \frac{1600\sqrt{13}}{46189} R_{60} + \frac{600\sqrt{17}}{4199} R_{80}$ $-\frac{10368\sqrt{21}}{96577} R_{10,0} + \frac{4356}{37145} R_{12,0} - \frac{264\sqrt{1001}}{37145} R_{12,4}$

Box matrix elements $P = (2\pi/L)(0, n, n)$, $S = \frac{1}{2}$

J'	L'	n'	J	L	n	$u_a^{-(L'+L+1)} B$
$\Lambda_B = G$ (partial)						
$\frac{5}{2}$	2	2	$\frac{9}{2}$	5	4	$-\frac{3\sqrt{105}}{308}iR_{30} - \frac{13\sqrt{14}}{924}iR_{32} - \frac{7\sqrt{165}}{286}iR_{50} + \frac{95\sqrt{154}}{3003}iR_{52}$ $-\frac{25\sqrt{462}}{2002}iR_{54} + \frac{915}{2288}iR_{70} + \frac{375\sqrt{21}}{16016}iR_{72}$ $-\frac{675\sqrt{462}}{16016}iR_{74} + \frac{15\sqrt{3003}}{2288}iR_{76}$
$\frac{5}{2}$	2	2	$\frac{9}{2}$	5	5	$-\frac{23\sqrt{30}}{924}R_{30} - \frac{95}{462}R_{32} - \frac{2\sqrt{2310}}{3003}R_{50} + \frac{2\sqrt{11}}{429}R_{52}$ $+\frac{16\sqrt{33}}{429}R_{54} + \frac{135\sqrt{14}}{2288}R_{70} + \frac{435\sqrt{6}}{2288}R_{72}$ $+\frac{105\sqrt{33}}{1144}R_{74} + \frac{45\sqrt{858}}{2288}R_{76}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	1	$\frac{\sqrt{105}}{13}R_{54} - \frac{\sqrt{105}}{65}R_{74} - \frac{\sqrt{2730}}{455}R_{76}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	2	$-\frac{5\sqrt{35}}{77}R_{32} + \frac{10\sqrt{385}}{1001}R_{52} - \frac{\sqrt{1155}}{1001}R_{54} - \frac{5\sqrt{210}}{2002}R_{72}$ $+\frac{2\sqrt{1155}}{715}R_{74} + \frac{3\sqrt{30030}}{1430}R_{76}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	3	$-\frac{5\sqrt{70}}{231}R_{30} + \frac{10\sqrt{21}}{231}R_{32} + \frac{10\sqrt{110}}{429}R_{50} + \frac{2\sqrt{231}}{273}R_{52}$ $-\frac{\sqrt{77}}{13}R_{54} - \frac{5\sqrt{6}}{143}R_{70} + \frac{27\sqrt{14}}{1001}R_{72} - \frac{3\sqrt{77}}{143}R_{74}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	4	$\frac{5\sqrt{7}}{11}R_{32} + \frac{8\sqrt{77}}{143}R_{52} - \frac{9\sqrt{231}}{1001}R_{54} - \frac{17\sqrt{42}}{286}R_{72}$ $-\frac{6\sqrt{231}}{1001}R_{74} - \frac{5\sqrt{6006}}{2002}R_{76}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	5	$\frac{5\sqrt{35}}{33}R_{30} + \frac{5\sqrt{42}}{231}R_{32} - \frac{7\sqrt{55}}{429}R_{50} - \frac{\sqrt{462}}{3003}R_{52}$ $+\frac{10\sqrt{154}}{1001}R_{54} - \frac{42\sqrt{3}}{143}R_{70} - \frac{6\sqrt{7}}{1001}R_{72} - \frac{15\sqrt{154}}{1001}R_{74}$
$\frac{5}{2}$	2	2	$\frac{11}{2}$	5	6	$\frac{50}{231}iR_{30} + \frac{5\sqrt{30}}{77}iR_{32} + \frac{5\sqrt{77}}{429}iR_{50} - \frac{3\sqrt{330}}{143}iR_{52}$ $+\frac{4\sqrt{105}}{715}iR_{70} - \frac{192\sqrt{5}}{715}iR_{72}$

Software overview

- C++ software: **BoxQuantization** class
- XML input to constructor (or use other structures)
 - specify total momentum \mathbf{d} , little group irrep Λ
 - dimensionless quantities $m_{\text{ref}}L$, ξ
 - for each channel:
 - masses m_{1a}/m_{ref} , m_{2a}/m_{ref}
 - particle spins s_{1a} s_{2a}
 - product of intrinsic parities $\eta_{1a}^P \eta_{2a}^P$
 - maximum orbital angular momentum $L_{\text{max}}^{(a)}$
 - if identical or not
- constructor automatically
 - sets up basis of states
 - constructs needed box matrices
 - constructs needed RGL zeta calculators
- for a given lab-frame E or E_{cm}
 - evaluates and returns \tilde{K} and/or $B^{(P)}$ matrices
 - evaluates and returns $[\det(1 - B^{(P)}\tilde{K})]^{1/N_{\text{det}}}$ or $[\det(\tilde{K}^{-1} - B^{(P)})]^{1/N_{\text{det}}}$
 - evaluates other quantities, too

Fitting (setting the stage)

- $\mathcal{E}(O)$ MC estimate of observable O using entire ensemble
- $\mathcal{E}_k^{(r)}(O)$ estimate of O from k -th resampling of scheme r
- common resampling schemes: jackknife $r = J$, bootstrap $r = B$
- covariance of O_i and O_j estimated using

$$\text{cov}(O_i, O_j) \approx \mathcal{N}^{(r)} \sum_{k=1}^{N_r} \left(\mathcal{E}_k^{(r)}(O_i) - \langle \mathcal{E}^{(r)}(O_i) \rangle \right) \times \left(\mathcal{E}_k^{(r)}(O_j) - \langle \mathcal{E}^{(r)}(O_j) \rangle \right),$$

$$\langle \mathcal{E}^{(r)}(O_i) \rangle = \frac{1}{N_r} \sum_{k=1}^{N_r} \mathcal{E}_k^{(r)}(O_i),$$

- for jackknife and bootstrap

$$\mathcal{N}^{(J)} = \frac{(N_J - 1)}{N_J}, \quad \mathcal{N}^{(B)} = \frac{1}{N_B - 1}$$

Fitting (still setting the stage)

- often goal is to describe set of observables by set of model functions containing unknown parameters
- arrange observables as components of vector \mathbf{R}
- fit parameters into a vector $\boldsymbol{\alpha}$
- denote the set of model functions by the vector $\mathbf{M}(\boldsymbol{\alpha}, \mathbf{R})$
- vector of residuals $\mathbf{r}(\mathbf{R}, \boldsymbol{\alpha}) = \mathbf{R} - \mathbf{M}(\boldsymbol{\alpha}, \mathbf{R})$
- in lattice QCD, get the best fit estimates by minimizing correlated $-\chi^2$ of residuals

$$\chi^2 = \mathcal{E}(r_i) \sigma_{ij}^{-1} \mathcal{E}(r_j),$$

- covariance $\sigma_{ij} = \text{cov}(r_i, r_j)$ of residuals
- best-fit parameter errors:
 - from minimization software
 - by minimizing $\chi_k^2 = \mathcal{E}_k^{(r)}(r_i) \sigma_{ij}^{-1} \mathcal{E}_k^{(r)}(r_j)$ for each resampling k and obtain covariances

Fitting subtleties

- if model **depends** on any observables, covariance matrix must be recomputed and inverted each time parameters α adjusted during minimization!
- if model **independent** of all observables $\text{cov}(r_i, r_j) = \text{cov}(R_i, R_j)$ simplifying minimization
- multiple ensembles
 - assume covariance zero between different ensembles, errors from minimization software, or
 - ensure N_r same for each ensemble, then apply above formulas
- primary goal here: best-fit estimates of κ_j parameters in \tilde{K} or \tilde{K}^{-1}
- two fitting methods follow

Fitting: spectrum method

- choose $E_{\text{cm},k}$ as observables
- model predictions by solving quantization for κ_j parameters
- problems:
 - root finding difficult, many computations of RGL zeta functions
 - ambiguity mapping model energies to observed energies
 - model predictions depend on observables m_{1a} , m_{2a} , L , ξ so MUST recompute covariance during minimization
- “Lagrange multiplier” trick removes obs. dependence in model
 - include m_{1a} , m_{2a} , L , ξ as both observables and model parameters
- observations

$$\text{Observations } R_i: \{ E_{\text{cm},k}^{(\text{obs})}, m_j^{(\text{obs})}, L^{(\text{obs})}, \xi^{(\text{obs})} \},$$

- model parameters

$$\text{Model fit parameters } \alpha_k: \{ \kappa_i, m_j^{(\text{model})}, L^{(\text{model})}, \xi^{(\text{model})} \},$$

Fitting: spectrum method (con't)

- residuals

$$r_k = \begin{cases} E_{\text{cm},k}^{(\text{obs})} - E_{\text{cm},k}^{(\text{model})}, & (k = 1, \dots, N_E), \\ m_{k'}^{(\text{obs})} - m_{k'}^{(\text{model})}, & (k = k' + N_E, k' = 1, \dots, N_p), \\ L^{(\text{obs})} - L^{(\text{model})}, & (k = N_E + N_p + 1), \\ \xi^{(\text{obs})} - \xi^{(\text{model})}, & (k = N_E + N_p + 2). \end{cases}$$

- compute $E_{\text{cm},k}^{(\text{model})}$ using only model parameters
- emphasize $E_{\text{cm},k}^{(\text{model})}$ very difficult to compute

Fitting: determinant residual method

- introduce quantization determinant as residual
- better to use function of matrix A with real parameter μ :

$$\Omega(\mu, A) \equiv \frac{\det(A)}{\det[(\mu^2 + AA^\dagger)^{1/2}]}$$

- model fit parameters are just κ_i parameters
- residuals

$$r_k = \Omega\left(\mu, 1 - B^{(P)}(E_{\text{cm},k}^{(\text{obs})}) \tilde{K}(E_{\text{cm},k}^{(\text{obs})})\right), \quad (k = 1, \dots, N_E),$$

- use only **observed** energies, particle masses, lattice size, anisotropy
- advantage: model predictions do not need root finding or RGL zeta computations
- model depends on observables, so covariance must be recomputed as κ_j parameters adjusted during minimization
- covariance recomputation still **much** simpler than root finding required in spectrum method

First test of fitting

- first test using determinant residual method
- applied to $I = 1 \rho \rightarrow \pi\pi$ system NPB 910, 842 (2016)
- included $L = 1$ and $L = 3$ partial waves
- fit forms:

$$(\tilde{K}^{-1})_{11} = \frac{6\pi E_{\text{cm}}}{g^2 m_\pi} \left(\frac{m_\rho^2}{m_\pi^2} - \frac{E_{\text{cm}}^2}{m_\pi^2} \right)$$

$$(\tilde{K}^{-1})_{33} = \frac{1}{m_\pi^7 a_3}$$

- results using only irreps in NPB 910, 842 (2016):


$$\frac{m_\rho}{m_\pi} = 3.351(25), \quad g = 5.96(26), \quad m_\pi^7 a_3 = -0.0015(27), \quad \chi^2/\text{dof} = 1.07$$

- results with additional $B_1^+ d^2 = 1$ irrep (no $L = 1$)

$$\frac{m_\rho}{m_\pi} = 3.353(24), \quad g = 6.00(26), \quad m_\pi^7 a_3 = -0.00029(240), \quad \chi^2/\text{dof} = 1.01$$

First test of fitting (con't)

- define Δ by $\tilde{K}_{11}^{-1} = B_{11}^{(P)}(1 + \Delta)$
- plot shows Δ consistent with zero in this energy range



Conclusion

- purpose: practitioner's guide plus software for carrying out two-particle scattering studies in lattice QCD
- quantization $\det(\tilde{K}^{-1} - B^{(P)}) = 0$, Hermitian “box matrix”
- provided explicit box matrix elements in block diagonal basis
 - several total momenta, spins $S \leq 2$, orbital $L \leq 6$
- software to include higher partial waves, multi-channels
- discussed two fitting strategies
- tested using $L = 1$ and $L = 3$ in $I = 1 \pi\pi$ scattering
- collaborators: John Bulava, Ben Hörz, Bijit Singha, Jacob Fallica, Drew Hanlon, Ruairi Brett
- related talks:
 - John Bulava: Pion-pion scattering 17:20 today
 - Ben Hörz: Multi-hadron spectroscopy 15:30 Thursday