

Development of Grassmann higher order tensor renormalization group

Yusuke Yoshimura, Yoshinobu Kuramashi¹, Yoshifumi Nakamura¹,
Shinji Takeda², Ryo Sakai²

Tsukuba Univ., RIKEN¹, Kanazawa Univ.²

June 21, 2017


35th International Symposium on Lattice Field Theory (Lattice 2017)
Palacio de Congresos de Granada

Introduction

- Tensor Renormalization Group (TRG) [M. Levin and C. P. Nave, Phys.Rev.Lett.99 (2007)]
 - Deterministic data compression
 - Completely free of the sign problem
- Outline of the TRG
 - Tensor network representation

$$Z = \sum_{\{l\}} \prod_n T_{l_{1,n} l_{1,n-1} l_{2,n} l_{2,n-2} l_{3,n} l_{3,n-3}}$$

- "Block Spin transformation" for the tensor network




- TRG study of relativistic QFTs on the lattice
 - 2D ϕ^4 model [Y. Shimizu, Chin.J.Phys.50 (2012)]
 - 2D Schwinger model [Y. Shimizu and Y. Kuramashi, Phys.Rev.D90 (2014); Phys.Rev.D90 (2014)]
 - 2D $N_f = 1$ Gross-Neveu model [S. Takeda and Y. Yoshimura, Prog.Theor.Exp.Phys.2015]
 - 3D free Wilson fermion system [R. Sakai and S. Takeda, Lattice 2016]

→ expectation, correlation function, second renormalization 2 / 15

- The procedure of HOTRG:

1. Insert "isometries" U, V into the network.
2. Contracting T, U , and V , obtain a new tensor T^{new} .



- If the bond dimension is D , the isometries can be considered $D^2 \times D_{\text{cut}}$ matrix ($D_{\text{cut}} \leq D^2$).
- D_{cut} is the keeping dimension for the truncation, and determines the accuracy of this approximation.
- If $D_{\text{cut}} = D^2$,

$$UU^\dagger = VV^\dagger = \text{identity}$$

- Computation time $\propto D_{\text{cut}}^{11}$, memory requirement $\propto D_{\text{cut}}^6$


Higher order SVD

- Isometries are made via higher order singular value decomposition.

$$M_{xx'yy'zz'} = \sum_{ijklmn} S_{ijklmn} U_{xi} U'_{x'j} V_{yk} V'_{y'l} W_{zm} W'_{z'n}$$

U, V, W, U', V', W' are unitary, and S possesses the orthogonality

$$\sum_{jklmn} S_{ijklmn} S'_{ijklmn} \propto \delta_{ii'}$$



For example:

$$\sum_{x'yy'zz'} M_{x_1x'yy'zz'}^* M_{x_2x'yy'zz'} = \sum_i U_{x_1i}^* \lambda_i U_{x_2i}$$

3D free Wilson fermion

- Wilson fermion action (Wilson parameter: $r = 1$)

$$S = \sum_{n,n'} \bar{\psi}_n D_{n,n'}^W \psi_{n'}, \quad D_{n,n'}^W = (m+3)\delta_{n,n'} - \frac{1}{2} \sum_{\mu,\pm} (1 \pm \gamma_\mu) \delta_{n,n' \pm \hat{\mu}}$$

- Gamma matrixes

$$\gamma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

- Partition function

$$Z = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \prod_n \left(\prod_s e^{-(m+3)\bar{\psi}_{n,s}\psi_{n,s}} \prod_\mu e^{\bar{\chi}_{\mu,n+\hat{\mu},1}\chi_{\mu,n,1}} e^{\bar{\chi}_{\mu,n,2}\chi_{\mu,n+\hat{\mu},2}} \right)$$

$$\chi_{1,n} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \psi_n, \quad \chi_{2,n} = \psi_n, \quad \chi_{3,n} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix} \psi_n$$

Tensor network representation

The procedure to transform into tensor network representation:

1. Expand each exponential binomially.
2. Insert new Grassmann numbers.
3. Separate original Grassmann numbers at other sites.
4. Integrate out original Grassmann numbers.

For example:

$$\begin{aligned} e^{\bar{\chi}_{\mu,n+\hat{\mu},1}\chi_{\mu,n,1}} &= \sum_{l_{\mu,n,1}=0}^1 (\bar{\chi}_{\mu,n+\hat{\mu},1}\chi_{\mu,n,1})^{l_{\mu,n,1}} \\ &= \sum_{l_{\mu,n,1}} \int (d\eta_{\mu,n,1}\eta_{\mu,n,1})^{l_{\mu,n,1}} (\bar{\chi}_{\mu,n+\hat{\mu},1}\chi_{\mu,n,1})^{l_{\mu,n,1}} \\ &= \sum_{l_{\mu,n,1}} \int (\chi_{\mu,n,1}d\eta_{\mu,n,1})^{l_{\mu,n,1}} (\bar{\chi}_{\mu,n+\hat{\mu},1}\eta_{\mu,n,1})^{l_{\mu,n,1}} \end{aligned}$$

- $\chi_{\mu,n,1}d\eta_{\mu,n,1}$ can be moved without -1 from the anticommutation.
- original $\chi_{\mu,n,1}$'s position is replaced by $\eta_{\mu,n,1}$.

Grassmann HOTRG

Corecting $\chi_{\mu,n,s} d\eta_{\mu,n,s}, \bar{\chi}_{\mu,n,s} d\bar{\eta}_{\mu,n,s}$, integrate out $\psi_n, \bar{\psi}_n$.

$$\mathcal{T}_{l_{1,n} l_{1,n-\hat{1}} l_{2,n} l_{2,n-\hat{2}} l_{3,n} l_{3,n-\hat{3}}} = T_{l_{1,n} l_{1,n-\hat{1}} l_{2,n} l_{2,n-\hat{2}} l_{3,n} l_{3,n-\hat{3}}}$$

$$\prod_{\mu} d\bar{\eta}_{\mu,n,2}^{l_{\mu,n,2}} d\eta_{\mu,n,1}^{l_{\mu,n,1}} d\eta_{\mu,n,2}^{l_{\mu,n-\hat{\mu},2}} d\bar{\eta}_{\mu,n,1}^{l_{\mu,n-\hat{\mu},1}} (\bar{\eta}_{\mu,n+\hat{\mu},1} \eta_{\mu,n,1})^{l_{\mu,n,1}} (\bar{\eta}_{\mu,n,2} \eta_{\mu,n+\hat{\mu},2})^{l_{\mu,n,2}}$$

- $T_{l_{1,n} l_{1,n-\hat{1}} l_{2,n} l_{2,n-\hat{2}} l_{3,n} l_{3,n-\hat{3}}}$ satisfies


$$\sum_{\mu,s} (l_{\mu,n,1} + l_{\mu,n-\hat{\mu},s}) \bmod 2 \neq 0 \Rightarrow T_{l_{1,n} l_{1,n-\hat{1}} l_{2,n} l_{2,n-\hat{2}} l_{3,n} l_{3,n-\hat{3}}} = 0$$

- Similarly, including out $\psi_n, \bar{\psi}_n$ for all sites,

$$Z = \sum_{\{l\}} \prod_n \mathcal{T}_{l_{1,n} l_{1,n-\hat{1}} l_{2,n} l_{2,n-\hat{2}} l_{3,n} l_{3,n-\hat{3}}}$$

- HOTRG procedure can be done like making tensor network representation.

Result: free energy



- The difference between our result and exact value

$$\delta = \frac{\ln Z_{\text{exact}} - \ln Z(D_{\text{cut}})}{\ln Z_{\text{exact}}}$$


- $V = 64^3$
- The accuracy increase as D_{cut} is enlarged.

Expectation value

- Impure tensor representation for $\langle \bar{\psi} \psi \rangle$:


$$\begin{aligned}\langle \bar{\psi}_{n_0} \psi_{n_0} \rangle &= \frac{1}{Z} \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \bar{\psi}_{n_0} \psi_{n_0} \\ &= \frac{1}{Z} \sum_{\{l\}} \mathcal{T}'_{n_0} \prod_{n \neq n_0} \mathcal{T}_n\end{aligned}$$

- Impure tensor contraction



- Isometries U, V are made from T .

Result: chiral condensate




- $V = 64^3$
- The accuracy almost unchanged as D_{cut} is enlarged.

Correlation function


- Impure tensor representation for $\langle \bar{\psi}_{n_1,1} \psi_{n_2,2} \rangle$:

$$\begin{aligned} \langle \bar{\psi}_{n_1,1} \psi_{n_2,2} \rangle &= \frac{1}{Z} \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \bar{\psi}_{n_1,1} \psi_{n_2,2} \\ &= \frac{1}{Z} \sum_{\{l\}} \mathcal{T}'_{n_1} \mathcal{T}''_{n_2} \prod_{n \neq n_1, n \neq n_2} \mathcal{T}_n \end{aligned}$$



- Basically, each impure tensor are renormalized independently.



- Someday, two impure tensors are combined one tensor.



Result: correlation function



- $V = 64^3, m = 1$
- The accuracy does not increase as D_{cut} is enlarged.
- At long distance, the accuracy decrease.

Second renormalization group [Xie et al. 2012]

- Second renormalization group (SRG) is the algorithm to take in global effects approximately.
- Environment tensor



$T^{(n)}$ is the tensor renormalized for n times.

- Density matrix


$$Z = \rho^{(n+1)} \cdot U^{(n+1)}$$

- singular value decomposition

$$\rho^{(n+1)} = u\sigma v^\dagger \Rightarrow U^{(n+1)} = vu^\dagger$$



Result: HOSRG



- $V = 64^3$
- The accuracy of HOSRG increase as compared with the accuracy of HOTRG.

Summary

- GHOTRG can calculate the chiral condensate and fermion correlator in 3D free Wilson fermion as impure tensor formalization.
- By the GHOSRG, the Free Energy accuracy increase.

In future→

- Applying GHOSRG scheme for tensor network including impure tensors to calculate expectation values and correlation functions.
- Verifying that GHOTRG is available in more complicated 3D model