

ChPT loops for the lattice: pion mass and decay constant; HVP at finite volume and $n\bar{n}$ -oscillations

ChPT loops
for the lattice

Johan Bijmens

Introduction

Two-point

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions



Johan Bijmens

Lund University



Vetenskapsrådet

bijmens@thep.lu.se
<http://thep.lu.se/~bijmens>
<http://thep.lu.se/~bijmens/chpt/>
<http://thep.lu.se/~bijmens/chiron/>

1 Introduction

2 Vector two-point functions for a_μ LO-HVP

- Connected and disconnected in infinite volume
- Finite volume
- Twisting
- Results

3 Pion mass and decay constant

4 $n\bar{n}$ oscillations

5 Conclusions

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point

Pion mass and
decay constant

$n\bar{n}$ oscillations


Conclusions

- ChPT = Effective field theory describing the lowest order pseudo-scalar representation
- or the (pseudo) Goldstone bosons from spontaneous breaking of chiral symmetry.
- The number of degrees of freedom depend on the case we look at
- Treat π, η, K as light and pointlike with a derivative and quark-mass expansion
- Recent review of LECs:

[JB, Ecker, Ann.Rev.Nucl.Part.Sci. 64 \(2014\) 149 \[arXiv:1405.6488\]](#)

Why

Muon: $a_\mu = (g - 2)/2$ and $a_\mu^{\text{LO,HVP}} = \int_0^\infty dQ^2 f(Q^2) \hat{\Pi}(Q^2)$



plot: $f(Q^2) \hat{\Pi}(Q^2)$ with $Q^2 = -q^2$ in GeV²

Figure and data: Aubin, Blum, Chau, Golterman, Peris, Tu,
Phys. Rev. D93 (2016) 054508 [arXiv:1512.07555]

Low energy quantity so ChPT should be useful

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point


(Dis)connected
Finite volume
Twisting
Results

Pion mass and
decay constant


$n\bar{n}$ oscillations

Conclusions

Two-point: Connected versus disconnected



Connected



Disconnected

yellow=lots of quarks/gluons

- $\Pi_{ab}^{\mu\nu}(q) \equiv i \int d^4x e^{iq \cdot x} \langle T(j_a^\mu(x) j_a^{\nu\dagger}(0)) \rangle$
- $j_{\pi^+}^\mu = \bar{d} \gamma^\mu u$
- $j_u^\mu = \bar{u} \gamma^\mu u$, $j_d^\mu = \bar{d} \gamma^\mu d$, $j_s^\mu = \bar{s} \gamma^\mu s$
- $j_e^\mu = \frac{2}{3} \bar{u} \gamma^\mu u - \frac{1}{3} \bar{d} \gamma^\mu d - \frac{1}{3} \bar{s} \gamma^\mu s$
- ChPT p^4 : Della Morte, Jüttner, JHEP 1011(2010)154 [arXiv:1009.3783]
- ChPT p^6 : JB, Relefors, JHEP 1611(2016)086 [arXiv:1609.01573]


Two-point: Connected versus disconnected

- Include also singlet part of the vector current
- There are new terms in the Lagrangian
- p^4 only one more: $\langle L_{\mu\nu} \rangle \langle L^{\mu\nu} \rangle + \langle R_{\mu\nu} \rangle \langle R^{\mu\nu} \rangle$
(drops out when subtracting $\Pi(0)$)
- \implies The pure singlet vector current does not couple to mesons until p^6
- \implies Loop diagrams involving the pure singlet vector current only appear at p^8 (implies relations)
- p^6 (no full classification, just some examples)
 $\langle D_\rho L_{\mu\nu} \rangle \langle D^\rho L^{\mu\nu} \rangle + \langle D_\rho R_{\mu\nu} \rangle \langle D^\rho R^{\mu\nu} \rangle,$
 $\langle L_{\mu\nu} \rangle \langle L^{\mu\nu} \chi^\dagger U \rangle + \langle R_{\mu\nu} \rangle \langle R^{\mu\nu} \chi U^\dagger \rangle, \dots$
- Results at two-loop order, unquenched isospin limit

Two-point: Connected versus disconnected


- $\Pi_{\pi^+\pi^+}^{\mu\nu}$: only connected
- $\Pi_{ud}^{\mu\nu}$: only disconnected
- $\Pi_{uu}^{\mu\nu} = \Pi_{\pi^+\pi^+}^{\mu\nu} + \Pi_{ud}^{\mu\nu}$
- $\Pi_{ee}^{\mu\nu} = \frac{5}{9}\Pi_{\pi^+\pi^+}^{\mu\nu} + \frac{1}{9}\Pi_{ud}^{\mu\nu}$
- Infinite volume (and the ab considered here):
$$\Pi_{ab}^{\mu\nu} = (q^\mu q^\nu - q^2 g^{\mu\nu}) \Pi_{ab}^{(1)}$$
- Large N_c + VMD estimate: $\Pi_{\pi^+\pi^+}^{(1)} = \frac{4F_\pi^2}{M_V^2 - q^2}$
- Plots on next pages are for $\Pi_{ab0}^{(1)}(q^2) = \Pi_{ab}^{(1)}(q^2) - \Pi_{ab}^{(1)}(0)$
- At p^4 the extra LEC cancels, at p^6 there are new LEC contributions, but no new ones in the loop parts

Two-point: Connected versus disconnected



- **Connected**
- p^6 is large
- Due to the L_i^r loops

Two-point: Connected versus disconnected




Disconnected

- p^6 is large
- Due to the L_i^r loops
- about $-\frac{1}{2}$ connected
- $-\frac{1}{10}$ is from

$$\Pi_{ee}^{(1)} = \frac{5}{9} \Pi_{\pi^+\pi^+}^{(1)} + \frac{1}{9} \Pi_{ud}^{(1)}$$

Two-point: with strange, electromagnetic current



- π connected u,d
 - ud disconnected u,d
 - ss strange current
 - us mixed $s-u,d$
 - new p^6 LEC cancels
 - Disconnected strange $\approx -15\%$ of total strange
- JB, Relefors,
LUTP 16-51 to appear

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point
(Dis)connected
Finite volume
Twisting
Results

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions

- One-loop calculation in finite volume done by
Aubin et al, Phys.Rev. D88 (2013) 7, 074505 [arXiv:1307.4701]
Aubin et al. Phys. Rev. D93 (2016) 054508 [arXiv:1512.07555]
and found to fit lattice data well
- two-loop in partially quenched
JB, Relefors, LU TP 16-51 to appear
- I will stay with ChPT and the p regime ($M_\pi L \gg 1$)
- $1/m_\pi = 1.4$ fm
may need to (and I will) go beyond leading $e^{-m_\pi L}$ terms
“around the world as often as you like”
- Convergence of ChPT is given by $1/m_\rho \approx 0.25$ fm

Finite volume and Twisted boundary conditions

- On a lattice at finite volume $p^i = 2\pi n^i/L$: very few momenta directly accessible
- Put a constraint on certain quark fields in some directions:
 $q(x^i + L) = e^{i\theta^i} q(x^i)$
- Then momenta are $p^i = \theta^i/L + 2\pi n^i/L$. Allows to map out momentum space on the lattice much better

Bedaque,...

- Small note:
 - Beware what people call momentum: is θ^i/L included or not?
 - Reason: a colour singlet gauge transformation $G_\mu^S \rightarrow G_\mu^S - \partial_\mu \epsilon(x)$, $q(x) \rightarrow e^{i\epsilon(x)} q(x)$, $\epsilon(x) = -\theta^i x^i/L$
 - Boundary condition
Twisted \Leftrightarrow constant background field+periodic

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point
(Dis)connected
Finite volume
Twisting
Results

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions

Drawbacks:

- Box: Rotation invariance \rightarrow cubic invariance
- Twisting: reduces symmetry further

Consequences:

- $m^2(\vec{p}^2) = E^2 - \vec{p}^2$ is not constant
- There are typically more form-factors
- In general: quantities depend on more (all) components of the momenta
- Charge conjugation involves a change in momentum

Two-point function: twisted boundary conditions

JB, Relfors, JHEP 05 (201)4 015 [arXiv:1402.1385]

- $\int_V \frac{d^d k}{(2\pi)^d} \frac{k_\mu}{k^2 - m^2} \neq 0$
- $\langle \bar{u} \gamma^\mu u \rangle \neq 0$
- $j_{\pi^+}^\mu = \bar{d} \gamma^\mu u$
satisfies $\partial_\mu \langle T(j_{\pi^+}^\mu(x) j_{\pi^+}^{\nu\dagger}(0)) \rangle = \delta^{(4)}(x) \langle \bar{d} \gamma^\nu d - \bar{u} \gamma^\nu u \rangle$
- $\Pi_a^{\mu\nu}(q) \equiv i \int d^4 x e^{iq \cdot x} \langle T(j_a^\mu(x) j_a^{\nu\dagger}(0)) \rangle$
Satisfies WT identity. $q_\mu \Pi_{\pi^+}^{\mu\nu} = \langle \bar{u} \gamma^\mu u - \bar{d} \gamma^\mu d \rangle$
- ChPT at one-loop satisfies this
see also Aubin et al, Phys.Rev. D88 (2013) 7, 074505 [arXiv:1307.4701]
- two-loop in partially quenched
JB, Relfors, LU TP 16-51, to appear
satisfies the WT identity (as it should)

ChPT loops
for the lattice

Johan Bijnens

Introduction


Two-point
(Dis)connected
Finite volume
Twisting
Results

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions

$$\langle \bar{u} \gamma^\mu u \rangle$$




Fully twisted

$\theta_u = (0, \theta_u, 0, 0)$, all others untwisted

$m_\pi L = 4$

For comparison: $\langle \bar{u} u \rangle^V \approx -2.4 \cdot 10^{-5} \text{ GeV}^3$
 $\langle \bar{u} u \rangle \approx -1.2 \cdot 10^{-2} \text{ GeV}^3$



Partially twisted

(ratio at $p^4 \equiv 2$ up to kaon loops)

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point

(Dis)connected

Finite volume

Twisting


Results

Pion mass and
decay constant


$n\bar{n}$ oscillations

Conclusions

Two-point partially twisted: components





Twisting spatially symmetric



Twisting in x-direction

- Small p^6 corrections (thin lines: p^4 only)
- $m_{\pi 0} L = 4 \quad m_{\pi 0} = 0.135 \text{ GeV}$
- $-q^2 \Pi_{\text{VMD}}^{(1)} = \frac{-4q^2 F_\pi^2}{M_V^2 - q^2} \approx 5e-3 \cdot \frac{q^2}{0.1} \implies$ **Correction at % level**
- Can use the difference between different twists with same q^2 to check the finite volume corrections

Two-point partially twisted: spatial average



- Plotted: volume correction to $\bar{\Pi} = \frac{1}{3} \sum_{i=x,y,z} \Pi^{ii}$
- Small p^6 corrections: compare left and right
- Can use the difference between different twists with same q^2 to check the finite volume corrections

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point

(Dis)connected

Finite volume

Twisting

Results

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions

Two flavour ChPT: mass and decay constant

- First step towards finding out why hard-pion ChPT does not work at three-loops
- Lowest order: Gell-Mann, Oakes, Renner (1968)
- Chiral logarithm Langacker, Pagels (1973)
- Full NLO (and properly starting ChPT) Gasser-Leutwyler (1984)
- NNLO Buergi (1996), JB, Colangelo, Ecker, Gasser, Sainio (1996)
- NNNLO JB, Hermansson-Truedsson (2017)

- LO and Chiral logs: current algebra
- NLO: Feynman diagrams (by hand) and direct expansion of functional integral (with REDUCE)
- NNLO: Feynman diagrams (with a little help from FORM)
- NNLO: Feynman diagrams purely with FORM
- Main stumbling block: integrals
 - Reduction to master integrals with REDUZE Studerus (2009)
 - Master Integrals known
Laporta-Remiddi (1996); Melnikov, van Ritbergen (2001)
- Lots of book-keeping: FORM
- Checks:
 - All nonlocal divergences must cancel
 - Use different parametrizations of the Lagrangian
 - Agree with known leading log result JB, Carloni (2009)

- LO and Chiral logs: current algebra
- NLO: Feynman diagrams (by hand) and direct expansion of functional integral (with REDUCE)
- NNLO: Feynman diagrams (with a little help from FORM)
- NNLO: Feynman diagrams purely with FORM
- Main stumbling block: integrals
 - Reduction to master integrals with REDUZE Studerus (2009)
 - Master Integrals known
Laporta-Remiddi (1996); Melnikov, van Ritbergen (2001)
- Lots of book-keeping: FORM
- Checks:
 - All nonlocal divergences must cancel
 - Use different parametrizations of the Lagrangian
 - Agree with known leading log result JB, Carloni (2009)

Diagrams



LUND
UNIVERSITY

ChPT loops
for the lattice

Johan Bijnens


Introduction

Two-point

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions



Results: LO or x -expansion / physical or ξ -expansion

- $x = \frac{M^2}{16\pi^2 F^2}, \quad L_x = \log \frac{M^2}{\mu^2}, \quad M^2 = 2B\hat{m}$

$$\frac{M_\pi^2}{M^2} = 1 + x \left(a_{11}^M L_x + a_{10}^M \right) + x^2 \left(a_{22}^M L_x^2 + a_{21}^M L_x + a_{20}^M \right) + x^3 \left(a_{33}^M L_x^3 + a_{32}^M L_x^2 + a_{31}^M L_x + a_{30}^M \right) + \dots$$

$$\frac{F_\pi}{F} = 1 + x \left(a_{11}^F L_x + a_{10}^F \right) + x^2 \left(a_{22}^F L_x^2 + a_{21}^F L_x + a_{20}^F \right) + x^3 \left(a_{33}^F L_x^3 + a_{32}^F L_x^2 + a_{31}^F L_x + a_{30}^F \right) + \dots$$

- $\xi = \frac{M_\pi^2}{16\pi^2 F_\pi^2}, \quad L_\pi = \log \frac{M_\pi^2}{\mu^2}$

- $\frac{M^2}{M_\pi^2} = 1 + \xi \left(b_{11}^M L_\pi + b_{10}^M \right) + \xi^2 \left(b_{22}^M L_\pi^2 + b_{21}^M L_\pi + b_{20}^M \right) + \xi^3 \left(b_{33}^M L_\pi^3 + b_{32}^M L_\pi^2 + b_{31}^M L_\pi + b_{30}^M \right) + \dots$

- $\frac{F}{F_\pi} = 1 + \xi \left(b_{11}^F L_\pi + b_{10}^F \right) + \xi^2 \left(b_{22}^F L_\pi^2 + b_{21}^F L_\pi + b_{20}^F \right) + \xi^3 \left(b_{33}^F L_\pi^3 + b_{32}^F L_\pi^2 + b_{31}^F L_\pi + b_{30}^F \right) + \dots$

$$\tilde{l}_i = 16\pi^2 l_i^r, \quad \tilde{c}_i = (16\pi^2)^2 c_i^r$$

a_{11}^M	$\frac{1}{2}$
a_{10}^M	$2\tilde{l}_3$
a_{22}^M	$\frac{17}{8}$
a_{21}^M	$-3\tilde{l}_3 - 8\tilde{l}_2 - 14\tilde{l}_1 - \frac{49}{12}$
a_{20}^M	$64\tilde{c}_{18} + 32\tilde{c}_{17} + 96\tilde{c}_{11} + 48\tilde{c}_{10} - 16\tilde{c}_9 - 32\tilde{c}_8 - 16\tilde{c}_7$ $- 32\tilde{c}_6 + \tilde{l}_3 + 2\tilde{l}_2 + \tilde{l}_1 + \frac{193}{96}$
a_{33}^M	$\frac{103}{24}$
a_{32}^M	$\frac{23}{2}\tilde{l}_3 - 11\tilde{l}_2 - 38\tilde{l}_1 - \frac{91}{24}$
a_{31}^M	$-416\tilde{c}_{18} - 208\tilde{c}_{17} - 32\tilde{c}_{16} + 96\tilde{c}_{14} + 8\tilde{c}_{13} - 48\tilde{c}_{12}$ $- 384\tilde{c}_{11} - 192\tilde{c}_{10} + 72\tilde{c}_9 + 144\tilde{c}_8 + 72\tilde{c}_7 + 64\tilde{c}_6 - 8\tilde{c}_5$ $- 56\tilde{c}_4 + 16\tilde{c}_3 + 32\tilde{c}_2 - 96\tilde{c}_1 - 8\tilde{l}_3^2 - 48\tilde{l}_3\tilde{l}_2 - 84\tilde{l}_3\tilde{l}_1$ $- \frac{88}{3}\tilde{l}_3 - \frac{231}{10}\tilde{l}_2 - \frac{69}{5}\tilde{l}_1 - \frac{74971}{8640}$
a_{30}^M	contains free p^8 LECs (and a lot more terms)

- Similar tables for a_i^F , b_i^M , b_i^F
- Coefficients depend on scale μ , but whole expression is μ -independent
- Can be rewritten in terms of scales in the logarithm rather than in terms of LECs à la FLAG
- Leading log: a number
- NLL: depends on l_i^r
- NNLL: depends on c_i^r
- For the mass all needed c_i^r can be had from mass, decay-constant and $\pi\pi$ parameters fitted to two-loop or p^6 (i.e. $r_M, r_F, r_1, \dots, r_6$).
- For decay need one more (busy checking if it can be had)

Results: numerics preliminary


ij	a_{ij}^M	b_{ij}^M	a_{ij}^F	b_{ij}^F
10	+0.00282	-0.00282	+1.09436	-1.09436
11	+0.5	-0.5	-1	+1
20	+1.65296	-1.65771	-0.04734	-1.15001
21	+2.4573	-3.29038	-1.90577	+4.13885
22	+2.125	-0.625	-1.25	-0.25
30	+0.39527	-6.7854	-244.499	242.236
31	-3.75977	+4.32719	-19.0601	32.1315
32	+17.1476	+0.62039	-9.39462	-6.77511
33	+4.29167	+5.14583	-3.45833	-0.41666

Note the large coefficients in the decay constant


Pion mass

$$F_\pi = 92.2 \text{ MeV}, F = F_\pi/1.037, \bar{l}_1 = -0.4, \bar{l}_2 = 4.3, \bar{l}_3 = 3.41, \bar{l}_4 = 4.51,$$

r_i from [JB et al 1997](#), other $c_i^r = 0$, $\mu = 0.77 \text{ GeV}$



x-expansion (F -fixed)



ξ -expansion (F_π fixed)

ξ -expansion converges notably better

ChPT loops
for the lattice

Johan Bijnens

Introduction


Two-point

Pion mass and
decay constant


$n\bar{n}$ oscillations

Conclusions

Pion decay constant



x-expansion (F -fixed)




ξ -expansion (F_π fixed)

- ξ -expansion converges better
- Large p^6 due to the “240” in a_{30}^F and b_{30}^F

- some GUT models have neutron-anti-neutron oscillations but no proton decay
- Limits:
 - $8.6 \cdot 10^7$ s from free neutrons (ILL)
 - $2.7 \cdot 10^8$ s from oxygen nuclei (super-K)
(\bar{n} mass inside nuclei very different from n -mass)
 - Possible ESS experiment improvement by up to 10^3

- Effective dimension 9 operator: “ $uuduud$ ”



- Classification of quark operators and RGE to two loops:
[Buchhoff, Wagman, Phys. Rev D93\(2016\)016005 \[arXiv:1506.00647\]](#)
and earlier papers in there
- [E. Kofoed, Master thesis LU TP 16-62; JB, Kofoed, in preparation](#)

- 14 operators
- Chiral representations under $SU(2)_L \times SU(2)_R$:


Chiral	#operators	Chiral	#operators
$(1_L, 3_R)$	3: Q_1, Q_2, Q_3	$(3_L, 1_R)$	3
$(5_L, 3_R)$	3: Q_5, Q_6, Q_7	$(3_L, 5_R)$	3
$(1_L, 7_R)$	1: Q_4	$(7_L, 1_R)$	1

- $n\bar{n}$ is $\Delta I = 1$ so bottom line needs isospin breaking
- For the others the different operators are different elements within the same representation
- Use heavy baryon formalism with a baryon (\mathcal{N}) and anti-baryon doublet (\mathcal{N}^c) with same velocity v
- These correspond two widely separated areas (v and $-v$) in the relativistic field: so no double counting

- $\mathcal{L} = \frac{F^2}{4} \langle u_\mu u^\mu + \chi_+ \rangle + \overline{\mathcal{N}} (i v \cdot D + g_A u \cdot S) \mathcal{N}$
 $+ \overline{\mathcal{N}^c} \tau^2 (i v \cdot D + g_A u \cdot S) \tau^2 \mathcal{N}^c + \text{higher orders}$
- $\mathcal{N} = \begin{pmatrix} p \\ n \end{pmatrix} \rightarrow h\mathcal{N}, \quad \overline{\mathcal{N}^c} = (\overline{p^c} \ \overline{n^c}) \rightarrow \overline{\mathcal{N}^c} h^T$
- $h(g_L, g_R, u)$: $SU(2)_V$ compensator chiral transformation
- Spurions for each quark $n\bar{n}$ operator:
 - $(1_L, 3_R)$: two $SU(2)_R$ doublet indices
 - $(5_L, 3_R)$: four $SU(2)_L$ and two $SU(2)_L$ doublet indices
 - $(1_L, 7_R)$: six $SU(2)_R$ double indices
 - plus parity conjugates


ChPT terms and diagrams

- p^0 :
 - $(1_L, 3_R)$: $(u\overline{\mathcal{N}^c})_{i_L} (u\mathcal{N})_{j_L}$
 - $(5_L, 3_R)$: $(u^\dagger\overline{\mathcal{N}^c})_{i_L} (u^\dagger\mathcal{N})_{j_L} (U_{T^2})_{k_R l_L} (U_{T^2})_{m_R n_L}$
 - $(7_L, 1_R)$: none (first one at p^2)
- p^1 : none that directly contribute (but in loops from p^3)
- p^2 : many (at least 20 each for $(3_L, 1_R)$ and $(3_L, 5_R)$)




- Q_1, Q_2, Q_3 same factor from loops (isospin)
- Q_5, Q_6, Q_7 same factor
(Conjecture: due to projection on $I = 1$ subspace)
- Q_1, Q_2, Q_3 :
$$1 + \frac{M_\pi^2}{16\pi^2 F_\pi^2} \left(\left(-\frac{3}{2} g_A^2 - 1 \right) \log \frac{M_\pi^2}{\mu^2} - 1 \right) + \text{order } p^2 \text{ LECs}$$
- Q_5, Q_6, Q_7 :
$$1 + \frac{M_\pi^2}{16\pi^2 F_\pi^2} \left(\left(-\frac{3}{2} g_A^2 - 7 \right) \log \frac{M_\pi^2}{\mu^2} - 1 \right) + \text{order } p^2 \text{ LECs}$$
- Also done at finite volume

Results (Preliminary)



Relative correction
from loops



Relative correction
from loops (absolute value)

Showed you results for:

- HVP: ChPT at two-loops including partially quenched
 - Connected versus disconnected at two-loops
 - Connected: twisting and finite volume at two-loops
- Two flavour ChPT correction at three loops for the pion mass and decay constant
- Two flavour ChPT correction at one-loop for $n\bar{n}$ -oscillations
- **Be careful: ChPT must exactly correspond to your lattice calculation**
- Programs available (for published ones) via CHIRON
Those for this talk: sometime later this year (I hope) (or ask me)

ChPT loops
for the lattice

Johan Bijnens

Introduction

Two-point

Pion mass and
decay constant

$n\bar{n}$ oscillations

Conclusions