


Solution of the sign problem in the Potts model at fixed baryon number

Urs Wenger

Albert Einstein Center for Fundamental Physics
University of Bern



in collaboration with

A. Alexandru, G. Bergner, D. Schaich

22 Juni 2017, Lattice 2017, Granada, Spain



Motivation: QCD in the heavy-dense limit

- ▶ Effective action in terms of (anti-)Polyakov loops P and P^\dagger .
- ▶ Canonical determinant $n_f = 3$ quarks:

$$\det \mathcal{D}_{n_f=3}^{HD} \sim 2 \sum_{\bar{x}} (\text{Tr } P_{\bar{x}})^2 \sum_{\bar{y}} \text{Tr } P_{\bar{y}} + 4 \sum_{\bar{x}} \text{Tr } P_{\bar{x}} \sum_{\bar{y} \neq \bar{x}} \text{Tr } P_{\bar{y}} \sum_{\bar{z}} \text{Tr } P_{\bar{z}}$$

- ▶ describes the propagation of baryons
- ▶ Invariant under global $\mathbb{Z}(3)$ -transformations
- ▶ Suffers from a severe sign problem, unless
 - ▶ all $P_{\bar{x}}$ align \iff deconfined phase
 - ▶ global $\mathbb{Z}(3)$ is promoted to a local one \iff strong coupling

- ▶ **Motivation**
 - ▶ QCD in the heavy-dense limit
 - ▶ Canonical formulation
 - ▶ Absence of the sign problem at strong coupling

- ▶ **Solution in the 3-state Potts model**
 - ▶ Canonical formulation
 - ▶ Bond formulation and cluster algorithm
 - ▶ Solution of the sign problem
 - ▶ cf. [Alford, Chandrasekharan, Cox, Wiese 2001]

- ▶ **Physics of the 3-state Potts model**
 - ▶ Density driven phase transition
 - ▶ (Anti-)Quark-(anti-)Quark correlators

The 3-state Potts model in $d = 3$ dimensions

- ▶ We use the **3-state Potts model** as a proxy for the effective Polyakov loop action of **heavy-dense QCD**.

- ▶ Grand-canonical partition function of the Potts model:

$$Z_{\text{GC}}(h) = \int \mathcal{D}z \exp(-S[z] + h \sum_x z_x)$$

- ▶ **Polyakov loops** are represented by the **Potts spins** $z_x \in \mathbb{Z}(3)$
- ▶ standard nearest-neighbour interaction

$$S[z] = -\gamma \sum_{\langle xy \rangle} \delta_{z_x, z_y}$$

- ▶ external 'magnetic' field $h = (2\kappa e^\mu)^\beta \Rightarrow$ breaks $\mathbb{Z}(3)$
- ▶ There is a **sign problem** for $h \neq 0$, i.e. at finite quark density.

The 3-state Potts model in $d = 3$ dimensions

- ▶ Canonical partition function for N_Q quarks:

$$Z_C(N_Q) = \sum_{\{n\}, |n|=N_Q} \int \mathcal{D}z \exp(-S[z]) \cdot \prod_x f[z_x, n_x]$$

- ▶ local quark occupation number $n_x \leq n_x^{\max}$ with $|n| = N_Q$
- ▶ use the simple local fermionic weights

$$f[z, n] = z^n$$

- ▶ equivalent to the grand-canonical partition function for small h
i.e. small density:

$$Z_{GC}(h) \simeq \sum_{N_Q=0}^{\infty} e^{\mu N_Q} Z_C(N_Q) \quad \text{for } h = e^{\mu} \ll 1$$

The 3-state Potts model in $d = 3$ dimensions

Canonical partition function

$$Z_C(N_Q) = \sum_{\{n\}} \int \mathcal{D}z \exp(\gamma \sum_{\langle xy \rangle} \delta_{z_x, z_y}) \prod_x z_x^{n_x}$$

- ▶ Action is manifestly complex \Rightarrow fermion sign problem!
- ▶ Global $\mathbb{Z}(3)$ symmetry ensures $Z_C(N_Q \neq 0 \bmod 3) = 0$:
 - ▶ projection onto integer baryon numbers
- ▶ In the limit $\gamma \rightarrow 0$, the global $\mathbb{Z}(3)$ becomes a local one:
 - ▶ projection onto integer baryon numbers on single sites

$$n_x = 0 \bmod 3 \quad (\text{limit } \gamma \rightarrow 0)$$

- ▶ sign problem is absent

Bond formulation and cluster algorithm

- ▶ Introduce bonds to express the action as

$$e^{\gamma \cdot \delta_{z_x, z_y}} = \sum_{b_{xy}=0}^1 (\delta_{z_x, z_y} \delta_{b_{xy}, 1} (e^\gamma - 1) + \delta_{b_{xy}, 0})$$

- ▶ The canonical partition function now becomes

$$Z_C(N_Q) = \sum_{\{n\}} \sum_{\{b\}} \int \mathcal{D}z \prod_{\langle xy \rangle} (\delta_{z_x, z_y} \delta_{b_{xy}, 1} (e^\gamma - 1) + \delta_{b_{xy}, 0}) \prod_x z_x^{n_x}$$

- ▶ sum over all bond configurations $\{b\}$

- ▶ Define the sum of bond weights over $\{n\}, \{b\}, \mathcal{D}z$ as $\langle\langle \cdot \rangle\rangle_{N_Q}$:

$$Z_C(N_Q) = \langle\langle \prod_x z_x^{n_x} \rangle\rangle_{N_Q}$$

Bond formulation and cluster algorithm

- ▶ $N_Q = 0$ gives the usual **Swendsen-Wang cluster construction**:
 - ▶ bond b_{xy} is occupied with probability $p(b_{xy} = 1) = (e^\gamma - 1)$, if $z_x = z_y$
 - ▶ weight of bond configuration is $W(\{b\}) = (e^\gamma - 1)^{N_b}$, with $N_b = \sum_{\langle xy \rangle} b_{xy}$
- ▶ Summation over $\mathbb{Z}(3)$ spins within connected cluster yields

$$Z_C(N_Q = 0) = \langle\langle 1 \rangle\rangle_{N_Q=0} = \sum_{\{b\}} (e^\gamma - 1)^{N_b} \cdot 3^{N_C}$$

- ▶ **total number of clusters N_C**
- ▶ cluster algorithm requires Euler-tour trees to achieve dynamic connectivity in $\mathcal{O}(\ln^2 V)$ instead of $\mathcal{O}(V \ln V)$ or $\mathcal{O}(V^2)$

Solution of the sign problem in the canonical formulation

- ▶ In the canonical formulation the cluster algorithm solves the sign problem:
 - ▶ include the fermionic contribution with an **improved estimator**
 - ▶ similar to idea in the grand canonical formulation

[Alford, Chandrasekharan, Cox, Wiese 2001]
- ▶ Average $\prod_x z_x^{n_x}$ over the subensemble of the 3^{N_C} configurations related by the $\mathbb{Z}(3)$ transformations:
 - ▶ total weight can be factorized into individual cluster weights $W_0(C)$,

$$\left[\prod_x z_x^{n_x} \right]_{3^{N_C}} = \left[\prod_C \prod_{x \in C} z_x^{n_x} \right]_{3^{N_C}} = \prod_C \left[\prod_{x \in C} z_x^{n_x} \right]_3 = \prod_C W_0(C)$$

where

$$W_0(C) = \left[\prod_{x \in C} z_x^{n_x} \right]_3 = \left[z^{\sum_{x \in C} n_x} \right]_3 = \begin{cases} 1 & \text{if } \sum_{x \in C} n_x = 0 \pmod{3}, \\ 0 & \text{else.} \end{cases}$$

Solution of the sign problem in the canonical formulation

- ▶ Hence, the canonical partition function becomes

Sign free canonical partition function

$$Z_C(N_Q) = \langle\langle \prod_x z_x^{n_x} \rangle\rangle_{N_Q} = \sum_{\{n\}} \sum_{\{b\}} (e^\gamma - 1)^{N_b} \cdot 3^{N_C} \cdot \prod_C \delta_{n_C, 0}$$

- ▶ $n_C = \sum_{x \in C} n_x \bmod 3$ denotes the **triality of the cluster C**
- ▶ $\prod_C \delta_{n_C, 0}$ projects on sector of configurations with triality-0 clusters only
- ▶ An intuitive, physical picture emerges:
 - ▶ only clusters with integer baryon number contribute
⇒ **confinement**
 - ▶ quarks can move freely within the cluster
⇒ **deconfinement within cluster**

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ First calculate weight for cluster C containing source z_y^* :

$$\left[\prod_{w \in C} z_w^{n_w - \delta_{w,y}} \right]_3 = \left[z^{\sum_{w \in C} n_w - \delta_{w,y}} \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 1 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 1}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right] \right]_3 = \left[\left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right] \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right] \right]_3 = \left[\left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right] \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

- ▶ Calculate the subensemble average including $z_x z_y^*$:

$$\begin{aligned} \left[\left[z_x z_y^* \prod_w z_w^{n_w} \right] \right]_{3^{N_C}} &= \delta_{C_x, C_y} \cdot \left[\left[z_x z_y^* \prod_w z_w^{n_w} \right] \right]_{3^{N_C}} \\ &+ (1 - \delta_{C_x, C_y}) \cdot \left[\left[z_x z_y^* \prod_w z_w^{n_w} \right] \right]_{3^{N_C}} \end{aligned}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right]_3 = \left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

- ▶ Calculate the subensemble average including $z_x z_y^*$:

$$\begin{aligned} \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} &= \delta_{C_x, C_y} \cdot \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} \\ &+ (1 - \delta_{C_x, C_y}) \cdot \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} \end{aligned}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right] \right]_3 = \left[\left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right] \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

- ▶ Calculate the subensemble average including $z_x z_y^*$:

$$\begin{aligned} \left[\left[z_x z_y^* \prod_w z_w^{n_w} \right] \right]_{3^{N_C}} &= \delta_{C_x, C_y} \cdot \prod_C \delta_{n_C, 0} \\ &+ (1 - \delta_{C_x, C_y}) \cdot \left[\left[z_x z_y^* \prod_w z_w^{n_w} \right] \right]_{3^{N_C}} \end{aligned}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right]_3 = \left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

- ▶ Calculate the subensemble average including $z_x z_y^*$:

$$\begin{aligned} \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} &= \delta_{C_x, C_y} \cdot \prod_C \delta_{n_C, 0} \\ &+ (1 - \delta_{C_x, C_y}) \cdot \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} \end{aligned}$$

Improved estimators for physical quantities

- ▶ We can define an improved estimator for the quark-antiquark correlator:

$$\langle z_x z_y^* \rangle_{N_Q} \equiv \langle\langle z_x z_y^* \prod_w z_w^{n_w} \rangle\rangle_{N_Q} / \langle\langle \prod_{w \in C} z_w^{n_w} \rangle\rangle_{N_Q}$$

- ▶ ... and the weight for cluster C containing source z_x :

$$\left[\prod_{w \in C} z_w^{n_w + \delta_{w,x}} \right]_3 = \left[z^{\sum_{w \in C} n_w + \delta_{w,x}} \right]_3 = \left\{ \begin{array}{ll} 1 & \text{if } \sum_{w \in C} n_w = 2 \pmod{3} \\ 0 & \text{else} \end{array} \right\} = \delta_{n_C, 2}$$

- ▶ Calculate the subensemble average including $z_x z_y^*$:

$$\begin{aligned} \left[z_x z_y^* \prod_w z_w^{n_w} \right]_{3^{N_C}} &= \delta_{C_x, C_y} \cdot \prod_C \delta_{n_C, 0} \\ &+ (1 - \delta_{C_x, C_y}) \cdot \delta_{n_{C_x}, 2} \cdot \delta_{n_{C_y}, 1} \prod_{\substack{C \neq C_x \\ C \neq C_y}} \delta_{n_C, 0} \end{aligned}$$

Improved estimators for physical quantities

- ▶ Similar expressions for $z_x z_y$, $z_x^* z_y^*$, z_x , $z_x^* = z_x^2$, ...
- ▶ An interesting quantity is the the **quark chemical potential**:

$$\mu(\rho) \equiv -\frac{1}{3} \log \frac{Z_C(N_Q + 3)}{Z_C(N_Q)}$$

- ▶ quark density $\rho \equiv (N_Q + 3/2)/V$
- ▶ partition function ratio can be computed using the multiplicity ratio average


$$\frac{Z_C(N_Q + 3)}{Z_C(N_Q)} = \left\langle \frac{\mathcal{M}(N_Q + 3, \{b\})}{\mathcal{M}(N_Q, \{b\})} \right\rangle_{N_Q}$$

and an estimator for the multiplicity ratio.

Physics of the 3-state Potts model

- Phase diagram in the $(e^\mu, \gamma) \equiv (h, \kappa)$ -plane:

[Alford, Chandrasekharan, Cox and Wiese 2001]




- deconfinement phase transition at $T = (0, 0.550565(10))$

Physics of the 3-state Potts model

- Phase diagram in the $(e^\mu, \gamma) \equiv (h, \kappa)$ -plane:

[Alford, Chandrasekharan, Cox and Wiese 2001]




- line of first order phase transitions from T to E

Physics of the 3-state Potts model

- Phase diagram in the $(e^\mu, \gamma) \equiv (h, \kappa)$ -plane:


[Alford, Chandrasekharan, Cox and Wiese 2001]



- critical endpoint $E = (0.000470(2), 0.549463(13))$

Physics of the 3-state Potts model


- ▶ Canonical simulation results in the deconfined phase:



- ▶ description in terms of a gas of (free) quarks

Physics of the 3-state Potts model


- ▶ Results from below the deconfinement transition:



- ▶ transition from the confined into the deconfined phase

Physics of the 3-state Potts model


- ▶ Results from below the deconfinement transition:



- ▶ typical signature of a 1st order phase transition

Physics of the 3-state Potts model


- ▶ Results from below the deconfinement transition:



- ▶ Maxwell construction yields critical μ_c

Physics of the 3-state Potts model


- ▶ Results from below the deconfinement transition:



- ▶ Maxwell construction yields critical μ_c

Physics of the 3-state Potts model


- ▶ Results from below the critical endpoint:



- ▶ crossover from the confined into the deconfined phase

Physics of the 3-state Potts model


- ▶ Results from below the critical endpoint:



- ▶ crossover from the confined into the deconfined phase

Physics of the 3-state Potts model


- ▶ (Anti)Quark-(anti)quark potentials at low temperature:



- ▶ confined phase: $\gamma = 0.3$ for $N_Q = 24$, $V = 16^3$, i.e. $\rho = 5.9 \cdot 10^{-3}$

Physics of the 3-state Potts model

- ▶ (Anti)Quark-(anti)quark potentials at low temperature:



- ▶ values at $r = 0$ and $r \rightarrow \infty$ match $\langle z \rangle, \langle z^* \rangle, \langle z^* \rangle \langle z^* \rangle, \dots$

Conclusions

- ▶ We solved the fermion sign problem for the $\mathbb{Z}(3)$ Potts model
 - ▶ isolate coherent dynamics of the $\mathbb{Z}(3)$ spins in clusters
 - ▶ cluster subaverages project on **nonzero, positive contributions**
⇒ increase of statistics exponential in N_C
- ▶ The solution provides an **appealing physical picture**:

Good algorithms reflect true physics insight! [Neuberger '17]

- ▶ quarks confined in clusters, but move freely within
 - ▶ at $\gamma \rightarrow 0$ clusters are confined to single sites only
 - ▶ **deconfinement** corresponds to appearance of a **percolating cluster**
- ▶ Extension to Polyakov loop models could be possible:
 - ▶ mechanism at work at $\beta = 0$
 - ▶ extend it to $\beta > 0$

Motivation: QCD in the heavy-dense limit

- ▶ Allows to describe **static quarks and anti-quarks**.
- ▶ Effective action in terms of **Polyakov loops P** and **anti-Polyakov loops P^\dagger** in $d = 3$:

Fermion determinant for $N_f = 2$ QCD

$$\det \mathcal{D}_a^{HD} = \prod_{\bar{x}} \det \left[\mathbb{I} + (2\kappa e^{+\mu})^{L_t} P_{\bar{x}} \right]^2 \det \left[\mathbb{I} + (2\kappa e^{-\mu})^{L_t} P_{\bar{x}}^\dagger \right]^2$$

- ▶ Canonical determinants obtained from transfer matrices.

Motivation: QCD in the heavy-dense limit

- ▶ Canonical determinant describing a **single quark**, i.e. $n_f = 1$:

$$\det \mathcal{D}_{n_f=1}^{HD} = \left((2\kappa)^{L_t} + (2\kappa)^{-L_t} \right) \cdot \sum_{\bar{x}} \text{Tr} P_{\bar{x}}$$

- ▶ For $n_f = 2$ quarks:

$$\begin{aligned} \det \mathcal{D}_{n_f=2}^{HD} / \Omega &\propto 2 \sum_{\bar{x}} \text{Tr} P_{\bar{x}} \sum_{\bar{y}} \text{Tr} P_{\bar{y}} \\ &+ \left(4 \sum_{\bar{x}} \text{Tr} P_{\bar{x}} \sum_{\bar{y}} \text{Tr} P_{\bar{y}} - 3 \sum_{\bar{x}} (\text{Tr} P_{\bar{x}})^2 + 2 \text{Tr} P_{\bar{x}}^\dagger \right) \end{aligned}$$

- ▶ Both determinants **vanish under global $\mathbb{Z}(3)$ -transformations**.

Motivation: QCD in the heavy-dense limit

- ▶ Canonical determinant $n_f = 3$ quarks:

$$\det \mathcal{D}_{n_f=3}^{HD} / \Omega = h_3 \cdot \left(4 \sum_{\bar{x}} \text{Tr} P_{\bar{x}}^\dagger \sum_{\bar{y}} \text{Tr} P_{\bar{y}} - 3 \sum_{\bar{x}} \text{Tr} P_{\bar{x}} \text{Tr} P_{\bar{x}}^\dagger + 2L_s^3 \right) \\ + h_1 \left(4 \sum_{\bar{x}} \text{Tr} P_{\bar{x}}^\dagger \sum_{\bar{y}} \text{Tr} P_{\bar{y}} + 2 \sum_{\bar{x}} (\text{Tr} P_{\bar{x}})^2 \sum_{\bar{y}} \text{Tr} P_{\bar{y}} \right. \\ \left. + 4 \sum_{\bar{x}} \text{Tr} P_{\bar{x}} \sum_{\bar{y} \neq \bar{x}} \text{Tr} P_{\bar{y}} \sum_{\bar{z}} \text{Tr} P_{\bar{z}} \right)$$

- ▶ describes the propagation of mesons and baryons
- ▶ Invariant under global $\mathbb{Z}(3)$ -transformations
- ▶ Suffers from a severe sign problem, unless
 - ▶ all $P_{\bar{x}}$ align \iff deconfined phase
 - ▶ global $\mathbb{Z}(3)$ is promoted to a local one \iff strong coupling