

# Chiral Extrapolations of the $\rho(770)$ Meson in $N_f = 2 + 1$ Lattice QCD Simulations

R. Molina<sup>1</sup>, B. Hu<sup>1</sup>, M. Doring<sup>1,2</sup>, M. Mai<sup>1</sup> and A. Alexandru<sup>1</sup>



(1)



(2)

# Contents

1. Unitarized Chiral Perturbation Theory
2. Summary of the results from analysis of  $N_f = 2$  lattice data
3. Analysis of the  $N_f = 2 + 1$  lattice data
4. Pion mass dependence of the  $\rho$  meson mass
5.  $N_f = 2$  vs.  $N_f = 2 + 1$  lattice data
6. Conclusions

# Unitarized Chiral Perturbation Theory

- **ChPT expansion of the amplitude for meson-meson scattering**

$$t(s) = t_2(s) + t_4(s) + \dots, t_{2k} = \mathcal{O}(p^{2k}) \quad (1)$$

- **Lowest-order Chiral Lagrangian:**

$$\mathcal{L}_2 = \frac{f^2}{4} \langle \partial_\mu U^\dagger \partial^\mu U + M(U + U^\dagger) \rangle \quad (2)$$

- **Next-to-leading order term:**

$$\begin{aligned} \mathcal{L}_4 = & L_1 \langle \partial_\mu U^\dagger \partial^\mu U \rangle^2 + L_2 \langle \partial_\mu U^\dagger \partial_\nu U \rangle \langle \partial^\mu U^\dagger \partial^\nu U \rangle \\ & + L_3 \langle \partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U \rangle + L_4 \langle \partial U^\dagger \partial^\mu U \rangle \langle U^\dagger M + M^\dagger U \rangle \\ & + L_5 \langle \partial_\mu U^\dagger \partial^\mu U (U^\dagger M + M^\dagger U) \rangle + L_6 \langle U^\dagger M + M^\dagger U \rangle^2 \\ & + L_7 \langle U^\dagger M - M^\dagger U \rangle^2 + L_8 \langle M^\dagger U M^\dagger U + U^\dagger M U^\dagger M \rangle \end{aligned} \quad (3)$$

# Unitarized Chiral Perturbation Theory

where  $U(\phi) = \exp(i\sqrt{2}\Phi/f)$ , and

$$\Phi(x) = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}}\eta \end{pmatrix} \quad (4)$$

$$M = \begin{pmatrix} m_\pi^2 & 0 & 0 \\ 0 & m_\pi^2 & 0 \\ 0 & 0 & 2m_K^2 - m_\pi^2 \end{pmatrix}. \quad (5)$$

J. Gasser and H. Leutwyler, *Annals Phys.* 158, 142 (1984)

J. Gasser and H. Leutwyler, *NPB250*, 465 (1985)

J. A. Oller, E. Oset and J. R. Pelaez, *PRD59*, 074001 (1999)

$T(\pi\pi \rightarrow \pi\pi)$  (1 channel) UChPT (NLO) Oller, Oset, Pelaez (1999)

$V_{\text{IAM}}$  is the kernel of the scattering equation:

$$T = T + VGT \longrightarrow T = [I - V_{\text{IAM}}G]^{-1}V_{\text{IAM}}, \text{ with}$$

$$V_{\text{IAM}} = [1 - V_4(V_2)^{-1}]^{-1}V_2$$

$$V_2 = \frac{m_\pi^2 - s}{f_\pi^2}$$

$$\begin{aligned} V_4 = & -\frac{4}{f^4}((2L_1 + L_3)(s - 2m_\pi^2)^2 \\ & + L_2((t - 2m_\pi^2)^2 + (u - 2m_\pi^2)^2) \\ & + 2(2L_4 + L_5)m_\pi^2(s - 2m_\pi^2) + \\ & 4(2L_6 + L_8)m_\pi^4) \end{aligned}$$

•  $I = 1, L = 1$ :

$$\longrightarrow V(\pi\pi) = -2p^2 / (3(f_\pi^2 - 8\hat{l}_1 m_\pi^2 + 4\hat{l}_2 E^2)) \quad (6)$$

and,  $\hat{l}_1 = 2L_4 + L_5$       $\hat{l}_2 = 2L_1 - L_2 + L_3$ .

Unitarized Chiral Perturbation Theory (Two channels:  $\pi\pi$ ,  $K\bar{K}$ )

- **Potential** ( $p$ -wave,  $I = 1$ ,  $\pi\pi$ ,  $K\bar{K}$ ):

$$V_2(W) = - \begin{pmatrix} \frac{2p_\pi^2}{3f_\pi^2} & \frac{\sqrt{2}p_K p_\pi}{3f_K f_\pi} \\ \frac{\sqrt{2}p_K p_\pi}{3f_K f_\pi} & \frac{p_K^2}{3f_K^2} \end{pmatrix}, \quad (7)$$

$$V_4(W) = - \begin{pmatrix} \frac{8p_\pi^2(2\hat{l}_1 m_\pi^2 - \hat{l}_2 W^2)}{3f_\pi^4} & \frac{8p_\pi p_K(L_5(m_K^2 + m_\pi^2) - L_3 W^2)}{3\sqrt{2}f_\pi^2 f_K^2} \\ \frac{8p_\pi p_K(L_5(m_K^2 + m_\pi^2) - L_3 W^2)}{3\sqrt{2}f_\pi^2 f_K^2} & \frac{4p_K^2(10\hat{l}_1 m_K^2 + 3(L_3 - 2\hat{l}_2)W^2)}{9f_K^4} \end{pmatrix}.$$

- **Scattering equation:**

$$T = V_2[V_2 - V_4 - V_2 G V_2]^{-1} V_2$$

$$S = \begin{pmatrix} \eta e^{2i\delta_1} & i(1 - \eta^2)^{1/2} e^{i(\delta_1 + \delta_2)} \\ i(1 - \eta^2)^{1/2} e^{i(\delta_1 + \delta_2)} & \eta e^{2i\delta_2} \end{pmatrix} \quad T_{ij} = -\frac{8\pi W}{2i\sqrt{p_i p_j}} (S_{ij} - \delta_{ij})$$

## Pion mass dependence of the meson decay constants

One loop SU(3) UChPT,

$$f_\pi = f_0 \left[ 1 - 2\mu_\pi - \mu_K + \frac{4M_{0\pi}^2}{f_0^2} (L_4^r + L_5^r) + \frac{8M_{0K}^2}{f_0^2} L_4^r \right]$$

$$f_K = f_0 \left[ 1 - \frac{3\mu_\pi}{4} - \frac{3\mu_K}{2} - \frac{3\mu_\eta}{4} + \frac{4M_{0\pi}^2}{f_0^2} L_4^r + \frac{4M_{0K}^2}{f_0^2} (2L_4^r + L_5^r) \right]$$

$$f_\eta = f_0 \left[ 1 - 3\mu_K + \frac{4L_4^r}{f_0^2} (M_{0\pi}^2 + 2M_{0K}^2) + \frac{4M_{0\eta}^2}{f_0^2} L_5^r \right]$$



J. Nebreda and J. R. Pelaez., Phys. Rev. D **81**, 054035 (2010)

Correlations in the energy levels:

$$\chi^2 = (\vec{W}_1 - \vec{W}_0)^T C^{-1} (\vec{W}_1 - \vec{W}_0)$$

$$W_{1i} = \frac{g(W_{0i}) - f(W_{0i})}{f'(W_{0i}) - g'(W_{0i})} + W_{0i} .$$

# Analysis of $N_f = 2$ lattice data: RQCD, GWU, QCDSF, Lang, ETMC and CP-PACS



$K\bar{K}$  channel included a posteriori (red)

B. Hu, R. Molina, M. Döring and A. Alexandru, PRL117 (2016)

G. S. Bali et al. [RQCD Collaboration], PRD93, 054509 (2016)

D. Guo, A. Alexandru, R. Molina and M. Doring, PRD94, 034501 (2016)


M. Göckeler et al. [QCDSF Collaboration], PoS LATTICE 2008, 136 (2008)

C. B. Lang, D. Mohler, S. Prelovsek and M. Vidmar, PRD84, 054503 (2011)

X. Feng, K. Jansen and D. B. Renner [ETMC Collaboration], PRD83, 094505 (2011)

S. Aoki et al. [CP-PACS Collaboration], PRD76, 094506 (2007)



# Analysis of $N_f = 2$ lattice data: RQCD, GWU, QCDSF, Lang, ETMC and CP-PACS



**Figure:** Left: Effect of the  $K\bar{K}$  channel in the  $(m_\rho, g)$  plane indicated with arrows, after chiral extrapolation to the physical pion mass. Right: error ellipses in the  $(\hat{l}_1, \hat{l}_2)$  plane.

Most ellipses in the  $(\hat{l}_1, \hat{l}_2)$  plane from the  $N_f = 2$  lattice data analysis have a common overlap area.

# $K\bar{K}$ loops in $\pi\pi \rightarrow \pi\pi$



Ratio of the couplings of the  $\rho$  meson to  $\pi\pi$  and  $K\bar{K}$

|                             | Oller/Pelaez1999 | Guo/Oller2012 |
|-----------------------------|------------------|---------------|
| $ g_{K\bar{K}}/g_{\pi\pi} $ | 0.54             | 0.64          |

NLO UChPT (PRD59) vs. One-loop UChPT (PRD84)

Not that small!

$N_f = 2 + 1$  lattice data analysis (UChPT)

- [1.] D. J. Wilson, R. A. Briceño, J. J. Dudek, R. G. Edwards and C. E. Thomas, Phys. Rev. D **92**, 094502 (2015). [Wilson15] [Str1-4]
- [2.] J. J. Dudek *et al.* [Hadron Spectrum Collaboration], Phys. Rev. D **87**, no. 3, 034505 (2013). [Dudek13] [Str1,3,4]
- [3.] J. Bulava, B. Fahy, B. Hörz, K. J. Juge, C. Morningstar and C. H. Wong, Nucl. Phys. B **910**, 842 (2016). [Bulava16] [Str1-3]
- [4.] J. Bulava, B. Hörz, B. Fahy, K. J. Juge, C. Morningstar and C. H. Wong, PoS LATTICE **2015**, 069 (2016). [Bulava15] [Str1-3]
- [5.] S. Aoki *et al.* [CS Collaboration], Phys. Rev. D **84**, 094505 (2011). [Aoki11] [Str1-3]
- [6.] C. Alexandrou *et al.*, arXiv:1704.05439 [hep-lat] (2017). [Alexandrou17] [Str3]
- [7.] Z. Fu and L. Wang, Phys. Rev. D **94**, no. 3, 034505 (2016). [Fu16] [Str3]
- [8.] T. Metivet [Budapest-Marseille-Wuppertal Collaboration], PoS LATTICE **2014**, 079 (2015). [BMW] (large errors-not analyzed)

## Minimization Strategies

- [Str1] Four parameters are left free:  $\hat{l}_1$ ,  $\hat{l}_2$ ,  $L_3$  and  $L_5$ ; Individual fits.
- [Str2]  $L_5$  is fixed. Combined fit of all data for  $m_\pi \leq 300$  MeV.
- [Str3]  $L_3$  and  $L_5$  are fixed. Individual fits.
- [Str4]  $L_3$  and  $L_5$  are fixed.<sup>1</sup> Combined fit of the Hadron Spectrum Collaboration data for  $m_\pi = 235$  and  $390$  MeV.

---

<sup>1</sup>Values from a fit to experimental phase shift data from  $\pi\pi$ ,  $\pi K$ ,  $K\bar{K}$ ..., M. Döring, U.-G. Meißner and W. Wang, JHEP **1310**, 011 (2013). “Experimental fit”

## Minimization strategy 1 (four free parameters)


Strong non trivial correlations, many  $\chi^2$  minima, best extrapolation to the physical point is selected.

|                |               | [Str1]               |                       |                         |                       |                          |
|----------------|---------------|----------------------|-----------------------|-------------------------|-----------------------|--------------------------|
|                | $M_\pi$ (MeV) | $\hat{I}_1$          | $\hat{I}_2$           | $L_3$                   | $L_5$                 | $\chi_{\text{d.o.f.}}^2$ |
| [Wilson15]     | 236           | $4.7^{+1.2}_{-0.8}$  | $-3.0^{+0.2}_{-0.9}$  | $-3.4^{+1.7}_{-0.2}$    | $-0.4^{+0.6}_{-1.0}$  | 1.0                      |
| [Dudek13]      | 391           | $5.4^{+0.9}_{-0.1}$  | $-7.4^{+0.6}_{-0.1}$  | $+4.3^{+0.3}_{-0.04}$   | $-3.8^{+0.7}_{-0.1}$  | 1.3                      |
| [Bulava16]     | 230           | $6.3^{+1.1}_{-1.0}$  | $-3.0^{+0.3}_{-0.4}$  | $-3.5^{+0.4}_{-0.3}$    | $-1.5^{+0.8}_{-1.0}$  | 1.3                      |
| [Bulava15]     | 280           | $6.3^{+1.8}_{-0.2}$  | $-3.0^{+0.5}_{-1.4}$  | $-3.0^{+3.3}_{-0.2}$    | $-0.4^{+1.0}_{-0.6}$  | 1.4                      |
| [Aoki11]       | 300 & 400     | $2.1^{+3.8}_{-1.2}$  | $-3.0^{+0.5}_{-1.9}$  | $-3.2^{+1.2}_{-0.6}$    | $0.7^{+0.7}_{-6.3}$   | 1.4                      |
| [Alexandrou17] | 317           | $13.3^{+0.8}_{-0.4}$ | $-12.9^{+0.5}_{-0.3}$ | $15.0^{+0.3}_{-0.1}$    | $-11.7^{+0.7}_{-0.4}$ | 0.7                      |
| [Str2]         | $\leq 300$    | $4.7^{+0.9}_{-0.5}$  | $-2.9^{+0.2}_{-0.1}$  | $-3.26^{+0.13}_{-0.06}$ | $0.64^{(f)}$          | 2.2                      |
| Experimental   | 138           | $0.26 \pm 0.05$      | $-3.96 \pm 0.04$      | $-3.01 \pm 0.02$        | $0.64 \pm 0.03$       | -                        |

**Table:** Low-energy constants and  $\chi_{\text{d.o.f.}}^2$  obtained in minimization strategy 1 and 2 for the  $N_f = 2 + 1$  lattice data analyzed.

A very good extrapolation in the UChPT model is possible, HOWEVER, some of the lattice data sets have VERY DIFFERENT values of the LECs. More constraints in the LECs needed.

## Minimization strategy 2 ( $L_5$ fixed)




**Figure:** Left: Phase shifts obtained for the minimization strategy 2. Lattice data included in the fit are [Wilson15], [Bulava16], [Bulava15], [Aoki11]. Right: Density histogram from resampling obtained in minimization 2. Ellipses in white show the result obtained in the fit of the  $N_f = 2$  data, [Lang11], with the SU(2) UChPT model. The star indicates the experimental fit.

## Minimization strategy 2 ( $L_5$ fixed)


- Strong correlations between  $\hat{l}_1$ ,  $\hat{l}_2$  and  $L_3$  are detected.
- In the common fit for  $m_\pi \leq 300$  MeV, [Aoki11] data are not well described.
- $\chi^2/\text{d.o.f} = 2.2$ : there are tensions between [Wilson15], [Bulava15/16] and [Aoki11].
- Differences with the values of the LECs from a fit to experimental phase shift data from  $\pi\pi$ ,  $K\bar{K}$ ,  $\pi K$ , ... ( $N_f = 2$  lattice data closer to the experimental point). Could be solved: 1) Fitting simultaneously lattice data from other reactions where these LECs are also involved. 2) Lattice determinations of  $f_\pi$ ,  $f_K$  for the lattice data analyzed. (New constraints)

## Minimization strategy 3 ( $L_3$ and $L_5$ fixed)



**Figure:** Left: Lattice data included in the fit are marked in red [Wilson15], [Dudek13], [Bulava16]. Right: Extrapolation to the physical point in comparison to the experimental data.

## Minimization strategy 3 ( $L_3$ and $L_5$ fixed)




**Figure:** Left: Lattice data included in the fit are marked in red [Bulava15], [Aoki11] and [Alexandrou17]. Right: Extrapolation to the physical point in comparison to the experimental data (red).

## Minimization strategy 3 ( $L_3$ and $L_5$ fixed)

|                       |               | [Str3]                                       |                  |                          |
|-----------------------|---------------|--|------------------|--------------------------|
|                       |               | ( $L_3 = -3.01^{(f)}$ , $L_5 = 0.64^{(f)}$ ) |                  |                          |
|                       | $M_\pi$ (MeV) | $\hat{l}_1$                                  | $\hat{l}_2$      | $\chi_{\text{d.o.f.}}^2$ |
| [Wilson15]            | 236           | $3.7 \pm 1.2$                                | $-3.2 \pm 0.3$   | 0.9                      |
| [Dudek13]             | 391           | $1.8 \pm 0.5$                                | $-3.7 \pm 0.3$   | 1.2                      |
| [Bulava16]            | 230           | $5 \pm 2$                                    | $-3.1 \pm 0.4$   | 1.1                      |
| [Bulava15]            | 280           | $5.7 \pm 1.3$                                | $-2.9 \pm 0.4$   | 1.2                      |
| [Aoki11]              | 300 & 400     | $2.5 \pm 0.7$                                | $-2.8 \pm 0.3$   | 1.1                      |
| [Alexandrou17]        | 317           | $2.8 \pm 0.8$                                | $-3.8 \pm 0.3$   | 0.6                      |
| [Str4]                | 236 & 391     | $2.0 \pm 0.2$                                | $-3.6 \pm 0.1$   | 1.1                      |
| [Guo16] ( $N_f = 2$ ) | 226 & 315     | $2.26 \pm 0.14$                              | $-3.44 \pm 0.03$ | 1.3                      |


**Table:** Low-energy constants and  $\chi_{\text{d.o.f.}}^2$  obtained in minimization strategies 3 and 4 for the  $N_f = 2 + 1$  lattice data. The superscript (f) indicates parameters held fixed at their experimental values. In the last row, the results from the analysis of the GWU  $N_f = 2$  lattice data are shown for comparison.

## Minimization strategy 3 ( $L_3$ and $L_5$ fixed)



**Figure:** Error ellipses (68% confidence) in the minimization 3 for the SU(3) analyses of the  $N_f = 2 + 1$  lattice data sets. The red dash-dotted ellipse (Lang11) represents the uncertainties from the analysis of the  $N_f = 2$  data using a one-channel SU(2) UChPT model.

## Minimization strategy 3




**Figure:**  $K\bar{K}$ -phase shifts (first row) and inelasticities (second row) obtained in the minimization strategy 3 for the different lattice data sets extrapolated at  $M_\pi = 236$  MeV, in comparison with the result from [Wilson15] (bands). In the bottom row, the extrapolated inelasticity to the physical point, in comparison with the experimental data (squares), and with the Roy-Steiner solution (black-dashed lines) is shown.

## Minimization strategy 3 ( $L_3$ and $L_5$ fixed)

- The minimization is stable compared to strategy 1.
- The values of the LECs are similar for the several  $N_f = 2 + 1$  data sets and similar to the ones found in  $N_f = 2$  analysis:
- $\hat{l}_2$  has a value of  $[-4 - (-3)] \times 10^{-3}$
- $\hat{l}_1$  is  $[2 - 4] \times 10^{-3}$  (except [Bulava15/16]  $[5 - 6] \times 10^{-3}$ )
- SU(2) and SU(3) analysis show similar results, indicating that the  $K\bar{K}$  dynamics can be absorbed in the SU(2) LECs for  $V(\pi\pi)$ .
- The extrapolated  $\rho$  mass of the [Dudek13] data is 25 MeV lighter ( $\sim 745$  MeV) than the experimental.
- The extrapolated  $\rho$  mass of the [Alexandrou17] data is 50 MeV lighter ( $\sim 720$  MeV).

## Minimization strategy 4 ( $L_3$ and $L_5$ fixed)

Combined fit of the Hadron Spectrum Col. data  $m_\pi = 236, 391$  MeV




**Figure:** Left: Lattice data included in the fit are marked in red [Wilson15] and [Dudek13]. Right: Extrapolation to the physical point in comparison to the experimental data.

$$\chi^2/\text{d.o.f} = 1.1$$

The extrapolated  $\rho$  mass is  $\sim 20$  MeV lighter ( $\sim 750$  MeV) than the experiment.


## Pion mass dependence of the $\rho$ meson mass



**Figure:** Pion mass dependence of the  $\rho$  meson mass as prediction of the combined fit of the data of [Wilson15] and [Dudek13] (red dotdashed upper band), in comparison with the result from the  $N_f = 2$  analysis of the GWU (blue continuous band and green curve, PRD94,034501), and with lattice data. The lower red band is the prediction in strategy (d) when the  $K\bar{K}$  channel is removed.

## Minimization strategy 4 ( $L_3$ and $L_5$ fixed)

- The pion mass dependence of the  $\rho$  mass obtained from a combined fit of the [Wilson15], [Dudek13] data is COMPATIBLE with most of the  $N_f = 2$  lattice data when the  $K\bar{K}$  channel is removed
- The analysis reveals tensions with [Bulava15/16] in the sense that these data do not predict a growing  $\rho$  mass with pion mass, as observed also in  $N_f = 2$  data
- From all the  $N_f = 2 + 1$  data, [Bulava15] ( $m_\pi = 280$  MeV) and [Alexandrou17] ( $m_\pi = 317$  MeV) are closer to the  $N_f = 2$  curve of  $m_\rho(m_\pi)$ .

Coupling,  $g$ , versus  $\rho$  mass

**Left:** minimization strategies 3 and 4; **Right:** 1 and 2. In the left figure, empty and filled symbols represent SU(2) and SU(3) analyses respectively. The experimental point is indicated as “phys.”. In the right figure, the red hexagon shows the result from minimization 2.

## $N_f = 2$ vs. $N_f = 2 + 1$ lattice data in the UChPT model

### $N_f = 2$ lattice data

- Have a common overlap area in the  $(\hat{l}_1, \hat{l}_2)$  plane
- The  $\rho$  mass clearly grows with the  $m_\pi$
- Most data fall in the same  $m_\rho(m_\pi)$  curve
- The coupling  $g$  is close to the experimental point
- The  $\rho$  mass is lighter than the experimental point ( $m_\rho \simeq 710$  MeV) for simulations close to the physical point [Bali] ( $m_\pi = 149$  MeV)


### $N_f = 2 + 1$ lattice data

- Do not have a common overlap area in the  $(\hat{l}_1, \hat{l}_2)$  plane
- The  $\rho$  mass do NOT GROW with  $m_\pi$  for [Bulava15/16], [Alexandrou17]. But It GROWS for [Wilson15][Dudek13]
- Some of the  $N_f = 2 + 1$  data are closer to the  $N_f = 2$   $m_\rho(m_\pi)$  curve ([Bulava15], [Alexandrou17])
- UChPT predicts a  $m_\rho \simeq 750$  MeV at the physical point from [Wilson15]/[Dudek13]

## Conclusions

- ★ We find tensions between most of the  $N_f = 2 + 1$  lattice data analyzed with the UChPT model, but NOT between  $N_f = 2$  data
- ★ However, there is mutual agreement between [Wilson15]/[Dudek13] data and  $N_f = 2$  lattice data when the  $K\bar{K}$  channel is removed/included, with  $\Delta m_\rho \simeq 40 - 50 \text{ MeV}$ , and close values of the LECs

## Fu16 data



**Figure:** First and third columns: phase shifts obtained in the SU(2) UChPT fits of the lattice data [Fu16]. The respective pion masses are indicated. Second and fourth columns: Respective chiral extrapolations to the physical point in comparison with the experimental data.

These data show discrepancies with other lattice data sets. In addition, they produce large values of the LECs in comparison with other data sets, and the UChPT extrapolation is far from the experiment.