

The spectrum of QCD at finite isospin density

Philipp Scior¹, Lorenz von Smekal²

¹Westfälische Wilhelms-Universität Münster

²Justus-Liebig-Universität Gießen

Lattice 2017



QCD at finite isospin chemical potential

- ▶ QCD with $N_f = 2$ and chemical potential

up quark: μ_u down quark: μ_d

- ▶ rewrite as baryon and isospin chemical potentials

$$\mu_B = \frac{3}{2}(\mu_u + \mu_d) \quad \mu_I = \mu_u - \mu_d$$



- ▶ μ_I introduces isospin imbalance
- ▶ occurs in nuclei, heavy-ion collisions, neutrons stars, possibly early universe
- ▶ here: $\mu_u = -\mu_d \rightarrow \mu_B = 0$
- ▶ fermion determinant: $\det(D(\mu_u)) \det(D(-\mu_u)) > 0 \rightarrow$ no sign problem

Phasediagram

Earlier works by:

- ▶ Son, Stephanov PRL86 (2001)
- ▶ Kogut, Sinclair PRD66 (2002)
- ▶ Detmold et al. PRD87 (2013)
- ▶ Kamikado et al. PLB718 (2013)
- ▶ Brandt, Enrodi, Gavai, ...

[this session]



Setup


- ▶ tree-level Symanzik improved gauge action
- ▶ finite pion source term

quark matrix:

$$M = \begin{pmatrix} D_\mu & \lambda\eta_5 \\ -\lambda\eta_5 & D_{-\mu} \end{pmatrix}$$

- ▶ D_μ is the staggered Dirac operator

Phasediagram



Lattice parameters:

$$\beta = 4.0, \quad m = 0.05, \quad \lambda = 0.005, \quad N_s = 16, \quad N_t = 24$$

Staggered Spectroscopy

- ▶ identify the operators corresponding to the desired states in the continuum limit

Operator	J^P	Particle
$\eta_4 \bar{\chi} \chi$	0^+	-
	0^{-+}	π
$\eta_i \eta_5 \zeta_i \bar{\chi} \chi$	1^{++}	a_1
	1^{--}	ρ
$\eta_i \eta_5 \eta_4 \zeta_i \zeta_4 \bar{\chi} \chi$	1^{+-}	b_1
	1^{--}	ρ
$\epsilon_{abc} \chi_a \chi_b \chi_c$	$\frac{1}{2}^-$	N(1535)
	$\frac{1}{2}^+$	N

Spectroscopy with finite pions source

- ▶ finite λ : quark propagator is non diagonal in isospin space
→ new terms in correlators

- ▶ disconnected contributions for all mesons $\sim \lambda^2$:

we choose $\frac{\lambda}{m} < 0.01 \rightarrow$ neglect disconnected contributions

- ▶ additional terms for some connected parts

$$\begin{aligned} \langle \pi^0(x) \bar{\pi}^0(0) \rangle = & \eta_4(x) \eta_5(x) \text{Tr}_c [G_{dd}^\dagger(x, 0, \mu) G_{uu}(x, 0, \mu) - G_{du}^\dagger(x, 0, \mu) G_{du}(x, 0, \mu) \\ & - G_{ud}^\dagger(x, 0, \mu) G_{ud}(x, 0, \mu) + G_{uu}^\dagger(x, 0, \mu) G_{dd}(x, 0, \mu)] \end{aligned}$$

- ▶ baryons: only connected pieces, additional terms

Spectroscopy at $\mu_I = 0$

two parametersets: both on $16^3 \times 24$ lattice:

(1) $\beta = 4.0$, $m = 0.05$, $\lambda = 0.005$

(2) $\beta = 3.5$, $m = 0.01$, $\lambda = 0.001$

Particle	(1)	(2)
am_π	0.66(1)	0.2597(1)
am_ρ	1.04(3)	-
am_N	1.64(5)	-


(1) pion is too heavy

(2) bad signal to noise ratio, unable to extract am_ρ , am_N

Spectroscopy at $\mu_l = 0$

π^+ correlator:


$$C(t) = A(-1)^t(e^{-m_\pi t} + e^{-m_\pi(N_t-t)})$$



Spectroscopy at finite μ_I


particles with positiv isospin are preferred over their anti-particles

$$C(t) = A(-1)^t (e^{-m_{\pi^+} t} + e^{-m_{\pi^-} + (N_t - t)})$$



Spectrum of finite μ_I

pion masses at finite μ_I in comparison with predictions from χ PT




Lattice parameters:

$$\beta = 4.0, \quad am = 0.05, \quad \lambda = 0.005, \quad N_s = 16, \quad N_t = 24$$

Spectrum of finite μ_l

pion masses at finite μ_l in comparison with predictions from χ PT




Lattice parameters:

$$\beta = 3.5, \quad am = 0.01, \quad \lambda = 0.001, \quad N_s = 16, \quad N_t = 24$$

Bad signal for all other hadron channels

Spectrum of finite μ_I

comparison of pion and ρ masses




Lattice parameters:

$$\beta = 4.0, \quad am = 0.05, \quad \lambda = 0.005, \quad N_s = 16, \quad N_t = 24$$

Spectrum of finite μ_I

baryon masses at finite μ_I

in comparison with predictions from χ PT



Summary & Conclusion

- ▶ QCD at finite isospin density
- ▶ started spectroscopy in- and outside pion condensation phase
- ▶ find better parameter set for simulations
- ▶ include disconnected contributions to meson correlators
- ▶ include stout-smearing
- ▶ $\lambda \rightarrow 0$ extrapolation