

# Double-winding Wilson loop in $SU(N)$ Yang-Mills theory

Ryutaro Matsudo

Graduate School of Science and Engineering, Chiba University

June 20, 2017

In collaboration with:

Kei-Ichi Kondo, Akihiro Shibata, Seikou Kato

arXiv:1706.05665


## Introduction

Recently, it is argued that **double-winding Wilson loops** can be used to test the candidates of the confinement mechanism in  $SU(2)$  gauge theory in Greensite and Höllwieser, Phys. Rev. D91, 054509 (2015).

In the paper, the following two types of models are considered:

- In the **center vortex model**, the expectation value of a double-winding Wilson loop is expected to decrease exponentially with **the difference of areas  $A_2 - A_1$** .
- In models associated with **the Abelian monopoles**, the expectation value is expected to decrease exponentially with **the sum of areas  $A_1 + A_2$** .


In lattice simulations, the expectation value of a double-winding Wilson loop decreases exponentially with **the difference of areas  $A_2 - A_1$**  in  $SU(2)$  Yang-Mills theory.




## Difference-of-areas behavior in center vortex model

We assume that the loop is so large that we can neglect the thickness of center vortices. Generally, if a vortex links the contour once, the phase factor is multiplied by  $-1$ , which is an element of the center of  $SU(2)$ .

- If a center vortex pierces  $A_1$ , then the vortex links the contour twice and hence the phase factor is multiplied by  $(-1)^2$ , which means there is no effect.
- If a center vortex pierces  $A_2 - A_1$ , then the vortex links the contour once and hence the phase factor is multiplied by  $-1$ .



$$U[C] \rightarrow (-1)^2 U[C] = U[C]$$



$$U[C] \rightarrow -U[C]$$

The situation is the same as if the contour is single winding and has the minimal area  $A_2 - A_1$ .  $\Rightarrow$  **difference-of-areas law**

## Sum-of-areas behavior in the models associated with Abelian monopole

The contour is rectangular.  $\Rightarrow W(C) \rightarrow \exp(-TV(R_1, R_2))$

In the dual superconductivity picture the electric flux between quark and antiquark forms a tube due to the condensation of the monopoles.


If two pairs are separated, the total energy is the sum of energy of each flux tube.

If  $\delta z \rightarrow 0$

TYPE I: The flux tubes attract each other and therefore the total energy is smaller than the separated case.


TYPE II: The flux tubes repel each other and therefore the total energy is almost sum of the energy of each flux.

In either case, the expectation value of double-winding Wilson loop is expected to decrease exponentially with either  $A_1$  and  $A_2$ .



## Purpose of this work

Our main purpose is to extend this argument to  $SU(N)$  case for  $N \geq 3$ . To do this we need to know the correct behavior of the expectation value of a double-winding Wilson loop in  $SU(N)$  Yang-Mills theory. In this talk, in the case that the two loops are identical we derive the behavior of the expectation value of the double-winding Wilson loop by using group identities and assuming the Casimir scaling irrespective of the dimensionality.



# Outline

- 1 Double-winding Wilson loop with identical loops
- 2  $N$ -times-winding Wilson loop
- 3 Double-winding Wilson loop with non-identical loops

# Outline

- 1 Double-winding Wilson loop with identical loops
- 2  $N$ -times-winding Wilson loop
- 3 Double-winding Wilson loop with non-identical loops

## Double-winding Wilson loop with identical loops

In  $SU(2)$  case, the expectation value of a double-winding Wilson loop with identical loops can be estimated by using the group identity

$$\text{tr} U^2 = \text{tr} U_A - 1,$$

where  $U$  is an arbitrary element of the group and  $U_A$  is the image of  $U$  under the adjoint representation.

Similarly, in  $SU(N)$  case, we can also estimate it by using the group identity

$$\text{tr} U^2 = \text{tr} U_{\square\square} - \text{tr} U_{\square},$$

where the Young diagrams denote the representations.

Using this group identity the double-winding Wilson loop with identical loops can be written as

$$W_2 = \frac{N+1}{2} W_{\square\square} - \frac{N-1}{2} W_{\square}.$$

## The expectation value

By using this relation and assuming **Casimir scaling**, we can estimate the expectation value for the intermediate size contour.

The ratios of the values of the quadratic Casimir in the representations and that in the fundamental representation are

$$C_2(\square\square) / C_2(F) = \frac{2(N+2)}{N+1},$$
$$C_2\left(\begin{array}{c} \square \\ \square \end{array}\right) / C_2(F) = \frac{2(N-2)}{N-1}.$$

By assuming the Casimir scaling, therefore, in the case that the minimal area  $S$  is intermediate, we can expect

$$\langle W_2 \rangle \simeq a_N \exp\left(-2\frac{N+2}{N+1}\sigma_F S\right) - b_N \exp\left(-2\frac{N-2}{N-1}\sigma_F S\right), \quad a_N, b_N > 0.$$

- For sufficiently large  $S$  the dominant part is the second term.
- For  $N \geq 3$ , this behavior is consistent with **neither difference-of-areas law nor sum-of-areas law**.

- In  $N = 3$  case, the behavior of the double-winding Wilson loop with identical loops is same as the single-winding one:

$$\langle W_2 \rangle \sim -b_3 e^{-\sigma_F S}.$$

- In large  $N$  limit, it is consistent with the sum-of-areas law:

$$\langle W_2 \rangle \sim -b_\infty e^{-2\sigma_F S}.$$

In the case that  $N$  is large, because the "string tension" approaches  $2\sigma_F$ , we need accurate calculation in order to distinguish the correct behavior from the behavior in the model associated with Abelian monopoles.

# Outline

- 1 Double-winding Wilson loop with identical loops
- 2  **$N$ -times-winding Wilson loop**
- 3 Double-winding Wilson loop with non-identical loops

## $N$ -times-winding Wilson loop

We can more easily distinguish the correct behavior from the behavior in models associated with Abelian monopoles by using  $N$ -times-winding Wilson loops. To see this, we consider the general  $m$ -times-winding Wilson loop with identical loops. In this case, we can also estimate the behavior of the expectation value by using the group identity.

## General $m$ -times-winding Wilson loop

The  $m$ -times-winding Wilson loop is defined as the Wilson loop for a contour which consists of  $m$  identical loops.

The trace of the  $m$ th power of a group element  $U$  can be written using the trace in the higher dimensional representation:

for  $m < N$

$$\mathrm{tr} U^m = U_{\square \cdots \square} - U_{\begin{array}{c} \square \\ \square \end{array} \cdots \square} + \cdots + (-1)^{l-1} U_{\begin{array}{c} \square \\ \vdots \\ \square \end{array} \cdots \square} + \cdots + (-1)^{m-1} U_{\begin{array}{c} \square \\ \vdots \\ \square \end{array}},$$

where all diagrams are "L-shape" or "I-shape", there are  $m$  boxes in each diagram, and there are  $l$  rows in the diagram in  $l$ th term, and for  $m \geq N$

$$\mathrm{tr} U^m = U_{\square \cdots \square} - U_{\begin{array}{c} \square \\ \square \end{array} \cdots \square} + \cdots + (-1)^{l-1} U_{\begin{array}{c} \square \\ \vdots \\ \square \end{array} \cdots \square} + \cdots + (-1)^{N-1} U_{\begin{array}{c} \square \\ \vdots \\ \square \end{array} \cdots \square},$$

where there are  $N$  terms.

Using this formula and assuming the Casimir scaling, in the case that the loop has an intermediate size, we can expect

$$\langle W_m \rangle \simeq \begin{cases} (-1)^{m-1} a_{Nm} \exp\left(-\frac{m(N-m)}{N-1} \sigma_F S\right) & \text{for } m < N, \\ (-1)^{N-1} a_{NN} & \text{for } m = N, \\ (-1)^{m-1} a_{Nm} \exp\left(-\frac{m(m-N)}{N+1} \sigma_F S\right) & \text{for } m > N. \end{cases}$$

The reason why, for  $m = N$ , the expectation value does not decrease exponentially is as follows. The young diagram




corresponds to the trivial representation if the number of the boxes is  $N$ . Hence, if  $m = N$ , the last term is equal to  $(-1)^{N-1}$ . In this way we see that  $\langle W_N \rangle$  does not follow the area law.

It seems that we can use  $W_N$  similarly to the double winding Wilson loop in  $SU(2)$  case.

## The behavior of the expectation value of $N$ -times-winding Wilson loop in the center vortex model and in models associated with Abelian monopoles

- If a center vortex pierces the minimal area of a  $N$ -times-winding Wilson loop, there is no effect because the  $N$ th power of an arbitrary element of the center is one. The expectation value is expected **not to decrease exponentially with the area**.
- Naively, in the models associated with monopole, the expectation value of  $N$ -times-winding Wilson loop is expected **to decrease exponentially with  $NS$** .



# Outline


- 1 Double-winding Wilson loop with identical loops
- 2  $N$ -times-winding Wilson loop
- 3 Double-winding Wilson loop with non-identical loops

## Double-winding Wilson loop with non-identical loops

I would like to conjecture the behavior of the double-winding Wilson loop average in the case that **two loops are non-identical**. To do this, we assume **factorization for non-intersecting coplanar loops**.


$$\langle \text{tr} U(C) \rangle = \langle \text{tr} U^2(C_1) U^{-1}(C_1) U(C_2) \rangle \simeq \text{tr} \langle U^2(C_1) \rangle \langle U^{-1}(C_1) U(C_2) \rangle$$

This assumption is consistent with usual area law.



By substituting the result in the identical loops case and usual single winding case, we obtain the dominant part as

$$\begin{aligned}
 W_2 &\sim -b_N \exp \left( -2 \frac{N-2}{N-1} \sigma_F A_1 - \sigma_F (A_2 - A_1) \right) \\
 &= -b_N \exp \left( -\sigma_F \left( \frac{N-3}{N-1} A_1 + A_2 \right) \right),
 \end{aligned}$$



which is consistent with  $A_1 = A_2$  case and  $A_1 = 0$  case.

According to Bralic (1980), in two-dimensional spacetime, we can calculate double-winding Wilson loops exactly as

$$\begin{aligned}
 W_2 &= \frac{N+1}{2} \exp \left( -\frac{g^2}{2} \frac{N^2-1}{2N} \left( \frac{N+3}{N+1} A_1 + A_2 \right) \right) \\
 &\quad - \frac{N-1}{2} \exp \left( -\frac{g^2}{2} \frac{N^2-1}{2N} \left( \frac{N-3}{N-1} A_1 + A_2 \right) \right).
 \end{aligned}$$

Therefore at least in two-dimensional spacetime, the above behavior is confirmed.

## Summary

- In the case that two loops are identical, we can expect the behavior of a double-winding Wilson loop expectation value by using the group identity and assuming the Casimir scaling in  $SU(N)$  case. It is consistent with **neither difference-of-areas law nor sum-of-areas law for  $N \geq 3$** . In large  $N$  limit, it is consistent with sum-of-areas law.
- We can also expect the behavior of a  $m$ -times-winding Wilson loop expectation value similarly.
- $N$ -times-winding Wilson loop expectation value in  $SU(N)$  gauge theory behaves similarly to the double-winding Wilson loop in  $SU(2)$  gauge theory.
- In the case that two loops are non-identical, we have conjectured the behavior of a double-winding Wilson loop expectation value, which is consistent with non-identical case. It is confirmed in two-dimensional Yang-Mills theory.

In the next talk, it is shown that a lattice simulation is consistent with the result of this talk for double-winding Wilson loop in  $SU(3)$  gauge theory.